

Transgression forms in dimension 4

by

Isabel M.C. Salavessa^{1*} and Ana Pereira do Vale^{2**}

Dedicated to Dmitri Alekseevsky on his 65th birthday

¹ Centro de Física das Interações Fundamentais, Instituto Superior Técnico, Edifício Ciência, Piso 3, 1049-001 LISBOA, Portugal; e-mail: isabel@cartan.ist.utl.pt

² Centro de Matemática, Universidade do Minho, Campus de Gualtar 4710-057 BRAGA, Portugal; e-mail: avale@math.uminho.pt

Abstract: We compute explicit transgression forms for the Euler and Pontrjagin classes of a Riemannian manifold M of dimension 4 under a conformal change of the metric, or a change to a Riemannian connection with torsion. These formulae describe the singular set of some connections with singularities on compact manifolds as a residue formula in terms of a polynomial of invariants. We give some applications for minimal submanifolds of Kähler manifolds. We also express the difference of the first Chern class of two almost complex structures, and in particular an obstruction to the existence of a homotopy between them, by a residue formula along the set of anti-complex points. Finally we take the first steps in the study of obstructions for two almost quaternionic-Hermitian structures on a manifold of dimension 8 to have homotopic fundamental forms or isomorphic twistor spaces.

1 Introduction

Some k -characteristic classes $Ch(E)$ of a vector bundle E over a manifold M , can be represented in the cohomology classes of M in terms of the curvature tensor defined with respect to a certain type of connections on E . If ∇, ∇' are such connections with curvature tensors R and R' , respectively, then Chern-Weil theory states that

$$Ch(R') = Ch(R) + dT$$

where T is a $(k-1)$ -form on M . A way to specify such a T is by pulling back each connection by the projection $\pi : M \times [0, 1] \rightarrow M$, $\pi(p, t) = p$, and then take the connection $\tilde{\nabla} = t\nabla' + (1-t)\nabla$ defined on $\pi^{-1}E$. Denoting by $\tilde{R}$ its curvature tensor, the Chern-Simons transgression $(k-1)$ -form on M , obtained by integration along $[0, 1]$ of the closed k -form $Ch(\tilde{R})$ on $M \times [0, 1]$,

$$T(\nabla, \nabla')(X_1, \dots, X_{k-1}) = \int_0^1 Ch(\tilde{R})\left(\frac{d}{dt}, X_1, \dots, X_{k-1}\right)dt$$

MSC 2000: Primary: 53C42, 53C55, 53C25, 53C38; Secondary: 57R20, 57R45.

Key Words: Characteristic classes, Transgression forms, Curvature tensors, almost complex structures, almost quaternionic Kähler, Singular connections, dimension 4 and 8.

*Partially supported by Fundação Ciência e Tecnologia through POCTI/MAT/60671/2004 and Plurianual of CFIF.

**Partially supported by Fundação Ciência e Tecnologia through POCTI/MAT/60671/2004.

satisfies $dT(\nabla, \nabla') = Ch(R') - Ch(R)$. To specify such T s is interesting by itself, but also when ∇' is a connection with singularities. This is the case when (M, g) is a Riemannian m -manifold with its Levi-Civita connection, and (E, g_E, ∇^E) is a Riemannian vector bundle of rank m such that there exists a conformal bundle map $\Phi : TM \rightarrow E$ which vanishes along a singular set Σ . This bundle map induces on $M \sim \Sigma$ a connection $\nabla' = \Phi^{-1*} \nabla^E$, that makes Φ a parallel bundle map. This connection can be seen as a singular Riemannian connection (with torsion) on M with respect to a degenerated metric $\hat{g}$ on M , but R' and $Ch(R')$ can be smoothly extended to Σ by the identities $R'(X, Y, Z, W) = g_E(R^E(X, Y)\Phi(Z), \Phi(W))$, $Ch(R') = Ch(R^E)$. If M is closed, and Ch gives a top rank form and an integral cohomology class, then $\int_M Ch(R)$ and $\int_M Ch(R')$ are finite integers, representing invariants. Moreover, if the singular set Σ is sufficiently small and regular, the Stokes theorem reads $\int_{M \sim V_\epsilon(\Sigma)} dT = - \int_{\partial V_\epsilon(\Sigma)} T$, where $V_\epsilon(\Sigma)$ is a tubular neighbourhood of Σ of radius ϵ , and letting $\epsilon \rightarrow 0$ may describe $Ch(E) - Ch(R)$ as a residue of T along Σ and expressed in terms of the zeros of Φ . This type of problem is studied in [15],[16],[17], using currents. We provide explicit formulae of transgression forms for the cases of the Euler and Pontrjagin classes. In section 10 we give some applications to minimal 4-submanifolds in Kähler-Einstein manifolds. In section 9 we describe an obstruction for two almost complex structures on M to be homotopic, measured by the difference between their Chern classes, translated to a residue formula on the set of anti-complex points. In section 11 we introduce the study of obstructions for two almost quaternionic-Hermitian structures on a Riemannian 8-manifold to have isomorphic twistor spaces or homotopic fundamental forms.

Let (M, g) be an oriented Riemannian manifold of dimension 4 with its Levi-Civita connection, curvature tensor R^M , Ricci tensor $Ricci^M$ and scalar curvature s^M . Our formulae are:

Proposition 1.1 *Let $f : M \rightarrow \mathbf{R}$ be a smooth map, and $\hat{g} = e^f g$. Set*

$$P(\nabla f) = (2\Delta f + \|\nabla f\|^2 - 2s^M)\nabla f + 4(Ricci^M)^\sharp(\nabla f) - \nabla(\|\nabla f\|^2). \quad (1.1)$$

Then

$$\begin{aligned} \mathcal{X}(\hat{R}^M) &= \mathcal{X}(R^M) + \frac{1}{32\pi^2} \text{div}_g(P(\nabla f)) \text{Vol}_M, \\ p_1(\hat{R}^M) &= p_1(R^M). \end{aligned} \quad (1.2)$$

The invariance of the Pontrjagin class under a conformal change of the metric is well known (see [6] or remark 3). We thank Sergiu Moroianu for drawing our attention to this. The authors do not recall to have seen formula (1.2) in the literature: $\text{div}_g(P(\nabla f))$ is a 2nd-order differential operator on ∇f . If $f = \log h$ where $h > 0$ except at a finite set of zeros p_α and poles p_i of homogeneous order $2k_\alpha$ and $2k_i$ respectively ($k_\alpha, k_i > 0$), then:

$$\frac{1}{32\pi^2} \int_M \text{div}_g(P(\nabla \log h)) \text{Vol}_M = \sum_\alpha -\frac{1}{2}k_\alpha^2(k_\alpha + 3) + \sum_i -\frac{1}{2}k_i^2(-k_i + 3).$$

Theorem 1.1 *Let $\Phi : TM \rightarrow E$ be a conformal bundle map into a Riemannian vector bundle (E, g_E, ∇^E) of rank 4 over M , with coefficient of conformality given by a non-negative function $h : M \rightarrow \mathbf{R}$ with zero set Σ . Let $\hat{g} = hg$, and $S' \in C^\infty(\otimes^2 T^*M \otimes TM)$ defined away from Σ :*

$$S'(X, Y) = \Phi^{-1}(\nabla_X \Phi(Y)) - \frac{1}{2}d \log h(X)Y - \frac{1}{2}d \log h(Y)X + \frac{1}{2}g(X, Y)\nabla \log h.$$

Then:

$$\begin{aligned}\mathcal{X}(R^E) &= \mathcal{X}(R^M) - \frac{1}{4\pi^2}d\left(\langle S' \wedge *(\hat{R}^M - \frac{1}{2}dS' - \frac{1}{3}(S')^2) \rangle_{\hat{g}}\right) + \frac{1}{32\pi^2}div_g(P(\nabla \log h))Vol_M \\ p_1(R^E) &= p_1(R^M) - \frac{1}{2\pi^2}d\left(\langle S' \wedge (\hat{R}^M - \frac{1}{2}dS' - \frac{1}{3}(S')^2) \rangle_{\hat{g}}\right)\end{aligned}$$

where $\hat{R}^M$ is the curvature tensor of $(M, \hat{g})$, and $S' : TM \rightarrow \Lambda^2 TM$, $(S')^2 : \Lambda^2 TM \rightarrow \Lambda^2 TM$, are defined by

$$\begin{aligned}\langle S'(X), Y \wedge Z \rangle_{\hat{g}} &= \hat{g}(S'(X, Y), Z) \\ \langle (S')^2(X \wedge Y), Z \wedge W \rangle_{\hat{g}} &= \hat{g}(S'(X, Z), S'(Y, W)) - \hat{g}(S'(X, W), S'(Y, Z)) \\ \langle \hat{R}^M(X \wedge Y), Z \wedge W \rangle_{\hat{g}} &= h\langle R^M(X \wedge Y) + \phi \bullet g(X \wedge Y), Z \wedge W \rangle_g \\ \phi &= \frac{1}{2}\left(-\frac{\|\nabla \log h\|^2}{4}g + \frac{1}{2}d \log h \otimes d \log h - Hess(\log h)\right).\end{aligned}$$

Moreover, if $d\Phi = 0$, then $\mathcal{X}(R^E) = \mathcal{X}(R^M) + \frac{1}{32\pi^2}div_g(P(\nabla \log h))Vol_M$ and $p_1(R^E) = p_1(R^M)$.

The angle $\theta \in [0, \pi]$ between two positive g -orthogonal almost complex structures J_0 and J_1 on M is defined by $\cos \theta = \frac{1}{4}\langle J_0, J_1 \rangle$, and $\Sigma = \{p \in M : J_1(p) = -J_0(p)\} = \cos \theta^{-1}(-1)$ is the set of *anti-complex points*. If J_1 is generic, Σ is a surface of M and an orientation can be given. Under the usual identification of J_0 with its Kähler form ω_0 , the orthogonal complement E_{J_0} of $\mathbf{R}J_0$ in $\Lambda_+^2 TM$, is a complex line bundle over M with complex structure " J_0 ". We denote by $\tilde{H}(p)$ the orthogonal projection of $J_1(p)$ into E_{J_0} . Let $N^1\Sigma$ be the total set of the unit normal bundle of Σ and $d_{N^1\Sigma}$ its Lebesgue measure. For each $(p, u) \in N^1\Sigma$ define $\kappa(p, u) \geq 1$ the order of the zero of $\phi_{(p, u)}(r) = (1 + \cos \theta)(exp_p(ru))$ at $r = 0$. We say that $(1 + \cos \theta)$ has a *controlled zero set* if there exist a non-negative integrable function $f : N^1\Sigma \rightarrow [0, +\infty]$ and $r_0 > 0$ s.t. $\sup_{0 < r < r_0} |r \frac{d}{dr} \log(\phi_{(p, u)}(r))| \leq f(p, u)$ a.e. $(p, u) \in N^1\Sigma$. For example, this holds if for all (p, u) , $\phi_{(p, u)}(r)$ is a polynomial function on r with coefficient of lowest order uniformly bounded away from zero. Some weaker conditions can be given on $\frac{d^k}{dr^k} \phi_{(p, u)}(0)$, $\frac{d^{k+1}}{dr^{k+1}} \phi_{(p, u)}(r)$ and $\frac{d^{k+2}}{dr^{k+2}} \phi_{(p, u)}(r)$, where $k = \kappa(p, u)$, to guarantee controlled zero set (see Prop. 9.4). For each $p \in \Sigma$, $S(p, 1)$ denotes the unit sphere of $T_p \Sigma^\perp$ and $\sigma_{d'}$ its volume. The function $\tilde{\kappa}(p) = \frac{1}{\sigma_{d'}} \int_{S(p, 1)} \kappa(p, u) d_{S(p, 1)} u$ is the *average order* of the zero p of $(1 + \cos \theta)$, in the normal direction. Assume M is compact.

Theorem 1.2 *If J_0 is almost Kähler then for any almost complex structure $J_1 = \cos \theta J_0 + \tilde{H}$*

$$\int_M (c_1(M, J_1) - c_1(M, J_0)) \wedge \omega_0 = \frac{1}{4\pi} \int_M div((\tilde{T}J_0)^\sharp) Vol_M \quad (1.3)$$

where $\tilde{T}$ is the 1-form on $M \sim \Sigma$, $\tilde{T}(X) = \frac{1}{(1+\cos \theta)} \langle \nabla_X \tilde{H}, J_0 \tilde{H} \rangle$. In the particular case that $\nabla^{E_{J_0}} \tilde{H}$ is J_0 -anti-complex we have

$$(1.3) = -\frac{1}{4\pi} \int_M \Delta \log(1 + \cos \theta) Vol_M. \quad (1.4)$$

In this case, assume Σ is a finite disjoint union of closed oriented submanifolds Σ^i of dimension $d_i \leq 2$. Let $\bigcup_\gamma k_{i\gamma}$ be the range set of κ on $N^1\Sigma^i$ and let $N^1\Sigma_\gamma^i = \kappa^{-1}(k_{i\gamma})$. If κ is bounded a.e.

and $(1 + \cos \theta)$ has a controlled zero set then a residue formula along Σ is obtained:

$$(1.3) = \frac{1}{2} \sum_{i:d_i=2} \int_{\Sigma^i} \tilde{\kappa}(p) Vol_{\Sigma^i} = \frac{1}{2} \sum_{i:d_i=2} \sum_{\gamma} k_{i\gamma} d_{N^1\Sigma^i}(N^1\Sigma_\gamma^i). \quad (1.5)$$

and so $\int_M \langle c_1(M, J_1), \omega_0 \rangle Vol_M \geq \int_M \langle c_1(M, J_0), \omega_0 \rangle Vol_M = \frac{1}{4\pi} \int_M (s^M + \frac{1}{2} \|\nabla \omega_0\|^2) Vol_M$,

with equality iff $d_i \leq 1 \ \forall i$. Thus, if $c_1(M, J_1) = c_1(M, J_0)$ (in $H^2(M, \mathbf{R})$) $\Sigma_i = \emptyset$ for $d_i = 2$.

Remark 1. Pairs of almost complex structures with or without the same properties may exist on a manifold. Alekseevsky [1] discovered examples of simply-connected non-compact Riemannian manifolds admitting a non-integrable almost-Kähler structure and also an integrable non-Kähler complex structure, namely some solvable groups of dimension $4(4 + p + q)$ ($q \neq 0$).

We also prove in section 9:

Proposition 1.2 *If M is compact, J_0 is Kähler and J_1 is almost Kähler then θ is constant. Thus, if $\cos \theta \neq \pm 1$, J_0 and J_1 are homotopic and define a hyper-Kähler structure on M .*

If (M^8, g) is an oriented 8-dimensional manifold and Q_0 and Q_1 are two almost quaternionic Hermitian structures (see [2] for definitions) we define an angle $\theta \in [0, \pi]$, by $\cos \theta = \frac{3}{10} \langle \Omega_0, \Omega_1 \rangle$, where Ω_i are the corresponding fundamental 4-forms. Let E_i be the corresponding rank 3 vector bundle generated by the twistor space of Q_i . In section 11 we prove:

Proposition 1.3 *If Q_0 is quaternionic Kähler and Q_1 is almost quaternionic Kähler and M is compact, then θ is constant. If $\cos \theta \neq -1$ then Ω_0 and Ω_1 are homotopic 4-forms in $H_+^4(M; \mathbf{R})$, and if Q_1 is also quaternionic Kähler, $p_1(E_0) = p_1(E_1)$. Furthermore, in the later case, (a), (b) or (c) must hold: (a) $E_0 = E_1$; (b) $E_0 \cap E_1$ has rank one, M is Kähler and both Q_0, Q_1 are locally hyper-Kähler structures; (c) $E_0 \cap E_1 = \{0\}$.*

We observe that the problem on a compact 4-manifold "almost Kähler + Einstein implies Kähler", also called the Goldberg-conjecture, is not completely solved. Salamon in [22] gives an example of a compact 8-dimensional almost quaternionic Kähler manifold that is not quaternionic Kähler. We thank the referee for drawing our attention to this reference.

2 Curvature tensors in dimension 4

Let V be a vector space of dimension 4 and with a inner product g . We identify $\Lambda^2 := \Lambda^2 V$ with $Skew(V)$ (the space of skew-symmetric endomorphisms of V) and with $\Lambda^2 V^*$ in the standard way, considering $Skew(V)$ as a subset of $V^* \otimes V$ with half of its usual Hilbert-Schmidt inner product. Consider the vector subspaces $\mathcal{R}$ and $\mathcal{R}^\perp$ of the symmetric and skew-symmetric linear endomorphisms of Λ^2 , respectively. $\mathcal{R}$, or more generally $L(\Lambda^2; L(V, V))$, is defined as the space of curvature tensors of V (see [7], [8], or [25] for details). We recall some definitions.

If $R \in L(\Lambda^2; L(V, V))$ let $\underline{R} \in L(V; L(\Lambda^2; V))$ given by $\underline{R}(Z)(X \wedge Y) := R(X \wedge Y)Z$. We also use the following notation: $R(X, Y, Z, W) = g(R(X \wedge Y)Z, W) = \underline{R}(Z, W, X, Y)$. If $R \in L(\Lambda^2; \Lambda^2)$, $\underline{R} = R^T$ (transposed). We assume that V is with a given orientation, and so the star operator $*$ $\in L(\Lambda^2; \Lambda^2)$ splits Λ^2 into its eigenspaces $\Lambda_\pm^2$, corresponding to the eigenvalues ± 1 , defining respectively the space of selfdual and of anti-self-dual two forms. Then

$R \in L(\Lambda^2; \Lambda^2)$ splits as $R = R_+^+ \oplus R_-^+ \oplus R_+^- \oplus R_-^-$ with $R_\pm^\pm \in L(\Lambda_\pm^2; \Lambda_\pm^2)$. The Ricci tensor $Ricci : L(\Lambda^2; \Lambda^2) \rightarrow L(V, V)$ and the scalar curvature are given respectively by $g(Ricci(R)(X), Y) = Ricci_R(X, Y) = tr\{Z \rightarrow R(X, Z)(Y)\}$ and $s_R = tr(Ricci_R) = 2\langle R, Id \rangle = 2tr(R)$. The sectional curvature of $R \in L(\Lambda^2; \Lambda^2)$ is denoted by $\sigma_R(P) = R(X, Y, X, Y)$ for each 2-plane P spanned by an o.n.b. $\{X, Y\}$, and the Bianchi map $b : L(\Lambda^2; L(V; V)) \rightarrow \Lambda^3 V^* \otimes V \subset L(\Lambda^2; L(V; V))$ is defined by $b(R)(X_1, X_2)(X_3) = R(X_1, X_2)(X_3) + R(X_3, X_1)(X_2) + R(X_2, X_3)(X_1)$. Then $b : \mathcal{R} \rightarrow \mathcal{R}, \mathcal{R}^\perp \rightarrow \mathcal{R}^\perp$. If $R \in \mathcal{R}$, $b(R) = \frac{1}{2}tr(*R)*$. Let $t(R) = \frac{1}{6}\langle b(R), * \rangle$. We denote by $Sym(V)$ the space of symmetric endomorphisms of V and by $Sym_0(V)$ its subspace of trace-free endomorphisms.

A complex 2-plane τ of V^c is said to be totally isotropic (t.i.) if $g(v, v) = 0 \forall v \in \tau$. If (e_i) is an o.n. basis and $\alpha = e_1 + ie_2, \beta = e_3 + ie_4$, then $\tau = span_{\mathbf{C}}\{\alpha, \beta\}$ defines a t.i. complex plane. Set $R(ijkl) = R(e_i, e_j, e_k, e_l)$. The isotropic sectional curvature w.r.t $R \in \mathcal{R}$ at τ is given by (see [21])

$$K_{isot}(R)(\tau) = \frac{R(\alpha \wedge \beta, \bar{\alpha} \wedge \bar{\beta})}{\|\alpha \wedge \beta\|^2} = \frac{1}{4} \left(R(1313) + R(1414) + R(2323) + R(2424) - 2R(1234) \right). \quad (2.1)$$

The *Kulkarni-Nomizu product* of $\xi, \phi \in \bigotimes^2 V^*$ is a symmetric product defined by

$$\xi \bullet \phi(X, Y, Z, W) = \xi(X, Z)\phi(Y, W) + \xi(Y, W)\phi(X, Z) - \xi(Y, Z)\phi(X, W) - \xi(X, W)\phi(Y, Z).$$

We have $Id = Id_{\Lambda^2} = \frac{1}{2}g \bullet g$. If $R \in L(\Lambda^2; L(V; V))$ it is defined the *Weitzenböck operator* $A(R)$

$$\langle A(R)(X, Y), Z \wedge W \rangle := Ricci_R \bullet g(X, Y, Z, W) + 2R(Z, X, Y, W) - 2R(W, X, Y, Z).$$

If $R \in \mathcal{R}$ then $A(R) \in \mathcal{R}$ and $A(R) = Ricci_R \bullet g - 2R + 2b(R)$. In this case, $Ricci_{A(R)} = s_R g$, $b(A(R)) = 4b(R)$. Furthermore, if $b(R) = 0$, $A(R) : \Lambda^\pm \rightarrow \Lambda^\pm$. Note that $* \in \mathcal{R}$, and if $R \in \mathcal{R}$, then $*R + R*, *R*$ still lie in $\mathcal{R}$ and $*R - R* \in \mathcal{R}^\perp$. Straightforward computations shows:

Proposition 2.1 *Let $R \in \mathcal{R}$ and $\tau = span_{\mathbf{C}}\{e_1 + ie_2, e_3 + ie_4\}$ with $e_1, \dots, e_4$ an o.n. basis with orientation ϵ . K_{isot} is computed at τ . Then*

$$\begin{array}{llll} Ricci_{Id} = 3g & s_{Id} = 12 & b(Id) = 0 & K_{isot}(Id) = 1 \\ Ricci_* = 0 & s_* = 0 & b(*) = 3* & K_{isot}(*) = -\frac{1}{2}\epsilon \\ Ricci_{*R*} = \frac{1}{2}s_R g - Ricci_R & s_{*R*} = s_R & b(*R*) = b(R) & K_{isot}(*R*) = K_{isot}(R) \\ Ricci_{R*} = Ricci_{*R} = t(R)g & s_{*R} = s_{R*} = 4t(R) & b(*R - R*) = 0 & K_{isot}(*R - R*) = 2(\sigma_R(34) - \sigma_R(12)). \end{array}$$

For $R, Q \in L(\Lambda^2; \Lambda^2)$, and $X, Y, Z, W \in V$ we use the following notation: $R(X \wedge Y) \wedge Q(Z \wedge W) = \langle R(X \wedge Y), (*Q)(Z \wedge W) \rangle Vol_V \in \Lambda^4 V^*$

Definition 2.1 *The Euler form $\mathcal{X}(R)$ and the Pontrjagin form $p_1(R)$ of $R \in L(\Lambda^2; \Lambda^2)$ are the 4-forms:*

$$\begin{aligned} 4\pi^2 \mathcal{X}(R) &= \underline{R}(e_1 \wedge e_2) \wedge \underline{R}(e_3 \wedge e_4) - \underline{R}(e_1 \wedge e_3) \wedge \underline{R}(e_2 \wedge e_4) + \underline{R}(e_1 \wedge e_4) \wedge \underline{R}(e_2 \wedge e_3) \\ &= \frac{1}{2} \langle R, *R* \rangle Vol_V = \frac{1}{2} \langle *R, R* \rangle Vol_V \end{aligned} \quad (2.2)$$

$$\begin{aligned} 4\pi^2 p_1(R) &= \underline{R}(e_1 \wedge e_2) \wedge \underline{R}(e_1 \wedge e_2) + \underline{R}(e_1 \wedge e_3) \wedge \underline{R}(e_1 \wedge e_3) + \underline{R}(e_1 \wedge e_4) \wedge \underline{R}(e_1 \wedge e_4) \\ &\quad + \underline{R}(e_2 \wedge e_3) \wedge \underline{R}(e_2 \wedge e_3) + \underline{R}(e_2 \wedge e_4) \wedge \underline{R}(e_2 \wedge e_4) + \underline{R}(e_3 \wedge e_4) \wedge \underline{R}(e_3 \wedge e_4) \\ &= \langle R^T, *R^T \rangle Vol_V = \langle R, R* \rangle Vol_V \end{aligned} \quad (2.3)$$

Note that $\langle R, Q \rangle = \langle R^T, Q^T \rangle$. Thus $\mathcal{X}(R^T) = \mathcal{X}(R)$, and $p_1(R^T) - p_1(R) = \langle R, *R - R* \rangle$.

A positive g -orthogonal complex structure J on V is a complex structure that induces the orientation of V and it is a linear isometry. Such structures are in 1-1 correspondence with the elements ω_J of $\bigwedge_+^2 V$ of norm $\sqrt{2}$, by $\omega_J(X, Y) = g(JX, Y)$. The condition of orthogonality between two of such J is equivalent to the anti-commuting condition, for the multiplication of complex structures corresponds to the quaternionic multiplication of two unit pure imaginary vectors of $\mathbf{R}^4$. For each J and $R \in L(\bigwedge^2; \bigwedge^2)$ we define the 2-forms on V

$$Ricci_{J,R}(X, Y) := \langle R(X \wedge Y), \omega_J \rangle = \underline{R}(\omega_J)(X, Y) \quad (2.4)$$

$$\Psi_J(R)(X, Y) := -\frac{1}{2}Ricci_R(X, JY) + \frac{1}{2}Ricci_R(Y, JX) \quad (2.5)$$

If $R \in \mathcal{R}$, then $\Psi_J(R)(X, Y) = Ricci^{(1,1)}(JX, Y)$. Usually $Ricci_{J,R}$ is denoted by $Ricci_*$ and named by *star-Ricci* but we do not use that notation to avoid confusion with the Ricci of the curvature tensor $*$. The *star-scalar* curvature is $s_{J,R} = 2\langle Ricci_{J,R}, \omega_J \rangle$, and $b_J(R) \in \bigwedge^2 V^*$ is defined by $b_J(R)(X, Y) = -\frac{1}{2}tr(Z \rightarrow Jb(R)(X, Y)(Z))$. If $P \in L_J(V; V)$, that is $P \circ J = J \circ P$ then both the complex trace $tr_J(P)$ and the complex determinant $det_J(P)$ are well defined. If $X_1, Y_1 = JX_1, X_2, Y_2 = JX_2$ is a real basis of V and for $\alpha = 1, 2$ denote " α " := $W_\alpha = \frac{1}{2}(X_\alpha - iY_\alpha)$, $\bar{\alpha} := W_{\bar{\alpha}} = \overline{W_\alpha}$, then $tr_J(P) = \sum_\alpha W_\alpha^* P^c(W_\alpha)$, $det_J(P) = det[W_\alpha^* P^c(W_\beta)]$ where $(W_\alpha^*, W_{\bar{\alpha}}^* = \overline{W_\alpha})$ is the complex dual basis of $\{W_\alpha, W_{\bar{\alpha}}\}$, and P^c denotes the complex linear extension of P to V^c . If $P \in L_J(V; V) \cap \bigwedge^2$, then $\langle P, \omega_J \rangle = tr_J(P)$. If $R \in L(\bigwedge^2; L_J(V; V))$, then R is said to be *J-invariant*.

Definition 2.2 If (V, J, g) is Hermitian, the first and second Chern form of $R \in L(\bigwedge^2; L_J(V; V))$ w.r.t. J , are respectively

$$c_1(R, J) = -\frac{i}{2\pi} Tr_J(\underline{R}) \quad c_2(R, J) = -\frac{1}{4\pi^2} det_J(\underline{R}). \quad (2.6)$$

It follows that, if $R \in L(\bigwedge^2; L_J(V; V) \cap \bigwedge^2)$ then $c_1(R, J) = \frac{1}{2\pi} Ricci_{J,R}$, $c_2(R, J) = \mathcal{X}(R)$, $p_1(R) = c_1(R, J) \wedge c_1(R, J) - 2c_2(R, J)$, and $b_J(R) = Ricci_{J,R} - \Psi_J$. If $R \in \mathcal{R}$ and is *J*-invariant, then $Ricci_R(JX, JY) = Ricci_R(X, Y)$, and $\Psi_J(R)(X, Y) = Ricci_R(JX, Y)$.

We denote by E_J the rank 2 subspace of $\bigwedge_+^2 V$ defined by the orthogonal complement of $\mathbf{R}\{\omega_J\}$

$$\bigwedge_+^2 V = \mathbf{R}\{\omega_J\} \oplus E_J, \quad \text{where } E_J = \{\omega \in \bigwedge_+^2 V : \omega(JX, JY) = -\omega(X, Y)\}$$

and a canonic complex structure can be given to E_J : $\tilde{J}\omega(X, Y) = -\omega(JX, Y)$. Let $\{e_1, e_2, e_3, e_4\}$ be a d.o.n.b. of V , giving a corresponding o.n.b. $\sqrt{2}\omega_\sigma$ of $\bigwedge_+^2 V$, $\omega_1 = e_1 \wedge e_2 + e_3 \wedge e_4$, $\omega_2 = e_1 \wedge e_3 - e_2 \wedge e_4$, $\omega_3 = e_1 \wedge e_4 + e_2 \wedge e_3$, and let J_σ be defined by $\omega_\sigma = g(J_\sigma(\cdot), \cdot)$. The 2-forms ω_2, ω_3 span E_J , and $\tilde{J}\omega_2 = \omega_3$ corresponds to $J_1 J_2 = J_3$. Any such o.n. (of norm $\sqrt{2}$) basis $(\omega_1, \omega_2, \omega_3)$, where $\omega_3 = "\omega_1 \omega_2"$ defines a canonic orientation on $\bigwedge_+^2 V$.

3 Almost Hermitian 4-manifolds

Assume (M, J, g) is an almost Hermitian 4-manifold with its Levi-Civita connection ∇ , and we use the above notation taking $V = T_p M$. (M, J, g) is said to be *almost Kähler* if the Kähler

form $\omega_J(X, Y) = g(JX, Y)$ is closed, and Kähler if ω_J is parallel. The latter is equivalent to J to be almost Kähler and integrable. Since $J^2 = -Id$ and J is g -orthogonal, $\forall X, Y, Z \in T_p M$

$$\nabla_Z J(JX) = -J\nabla_Z J(X), \quad g(\nabla_Z J(X), Y) = -g(X, \nabla_Z J(Y)). \quad (3.1)$$

We use the following sign for curvature tensors $R(X, Y) = -\nabla_X \nabla_Y + \nabla_Y \nabla_X + \nabla_{[X, Y]}$, $\forall X, Y \in T_p M$, and denote by R^M the curvature tensor of M , $Ricci_J^M = Ricci_{J, R^M}$, $s_J^M = s_{J, R^M}$, and by $\langle \cdot, \cdot \rangle$ the Hilbert Schmidt inner product on $\wedge^k TM^*$. The Weitzenböck formulae for ω_J read (see e.g. [8])

$$\Delta \omega_J = -tr \nabla^2 \omega_J + A(R^M)(\omega_J) \quad (3.2)$$

$$0 = \frac{1}{2} \Delta^+ \|\omega_J\|^2 = -\langle \Delta \omega_J, \omega_J \rangle + \|\nabla \omega_J\|^2 + \langle A(R^M) \omega_J, \omega_J \rangle. \quad (3.3)$$

Since ω_J is a self-dual form, $\|d\omega_J\| = \|\delta\omega_J\|$. Thus J is almost Kähler iff $\delta\omega_J = 0$. In this case ω_J is harmonic. If J is not Kähler we may use the *canonical Hermitian* connection (see [13])

$$\tilde{\nabla}_X Y = \nabla_X Y - \frac{1}{2} J(\nabla_X J(Y)).$$

This connection satisfies $\tilde{\nabla} g = \tilde{\nabla} J = 0$, but has torsion $\tilde{T}(X, Y) = -\frac{1}{2} J dJ(X, Y)$. This is a $U(2)$ -connection on M , and so its curvature tensor $\tilde{R}$ is J -invariant. The Chern classes of M using $\tilde{R}$ satisfy $2\pi c_1(M, J) = Ricci_{J, \tilde{R}}$, and a direct computation shows that

$$Ricci_{J, \tilde{R}}(X, Y) = Ricci_J^M(X, Y) + \eta_J(X, Y), \quad s_{J, \tilde{R}} = s_J^M + 2\langle \eta_J, \omega_J \rangle$$

where η_J is the 2-form on M

$$\eta_J(X, Y) = \frac{1}{4} \langle J \nabla_X J, \nabla_Y J \rangle, \quad (3.4)$$

and the inner product is the Hilbert-Schmidt inner product on $TM^* \otimes TM$. If (M, J, g) is almost Kähler then (see [5] for a survey)

$$\nabla_{JX} J = -J(\nabla_X J). \quad (3.5)$$

In this case $\tilde{T}^{(1,1)} = 0$, and $\langle \eta_J, \omega_J \rangle = -\frac{1}{8} \|\nabla J\|^2$. Furthermore, for almost Kähler J (see [9])

$$s_J^M - s^M = \|\nabla \omega_J\|^2 \quad (3.6)$$

$$4\pi \langle c_1(M, J), \omega_J \rangle = 2\langle \tilde{R}(\omega_J), \omega_J \rangle = s^M + \frac{1}{2} \|\nabla \omega_J\|^2 = \frac{1}{2} (s_J^M + s^M). \quad (3.7)$$

We consider on E_J the induced connection ∇^{E_J} from the connection ∇^+ of $\wedge_+^2$.

Proposition 3.1 *If (M, J, g) is almost Hermitian*

$$\begin{aligned} 2\pi c_1(E_J)(X, Y) &= Ricci_J^M(X, Y) + \eta_J(X, Y) = 2\pi c_1(M, J) \\ p_1(\wedge_+^2 T_p M) &= c_1(E_J) \wedge c_1(E_J). \end{aligned}$$

Proof. Let $\frac{\omega_2}{\sqrt{2}}, \frac{\omega_3}{\sqrt{2}}$ be a local d.o.n. frame of E_J , and $(\cdot)^{E_J}$ denote the orthogonal projection onto E_J . Since $\|\omega_J\| = \sqrt{2}$ is constant, $\nabla_Y^+ \omega_J$ is a section of E_J . Now,

$$\begin{aligned} \nabla_Y^{E_J} \nabla_X^{E_J} s &= (\nabla_Y^+ \nabla_X^{E_J} s)^{E_J} = (\nabla_Y^+ \nabla_X^+ s - \frac{1}{2} \langle \nabla_X^+ s, \omega_J \rangle \nabla_Y^+ \omega_J)^{E_J} \\ &= (\nabla_Y^+ \nabla_X^+ s)^{E_J} + \frac{1}{2} \langle s, \nabla_X^+ \omega_J \rangle \nabla_Y^+ \omega_J. \end{aligned}$$

Thus

$$R^{E_J}(X, Y)s = (R^+(X, Y)s)^E + \frac{1}{2}\langle s, \nabla_X^+ \omega_J \rangle \nabla_Y^+ \omega_J - \frac{1}{2}\langle s, \nabla_Y^+ \omega_J \rangle \nabla_X^+ \omega_J. \quad (3.8)$$

The curvature tensor of Λ_+^2 satisfies $\langle R^+(X, Y)\omega_2, \omega_3 \rangle = 2R^M(X, Y)\omega_J$. E_J is a complex line bundle over M and so it has a (real) volume element Vol_{E_J} . Using (3.1) we easily see that another way to express η_J is $\eta_J(X, Y) = \frac{1}{2}Vol_{E_J}(\nabla_X^+ \omega_J, \nabla_Y^+ \omega_J)$. Then

$$2\pi c_1(E_J)(X, Y) := \langle R^{E_J}(X, Y) \frac{\omega_2}{\sqrt{2}}, \frac{\omega_3}{\sqrt{2}} \rangle = R^M(X, Y)\omega_J + \eta_J(X, Y). \quad (3.9)$$

Since $c_2(\mathcal{C}\{\omega_J\} \oplus E_J^c) = c_1(\mathcal{C}\{\omega_J\}) \wedge c_1(E_J^c) - c_2(E_J^c)$ and $c_1(E_J^c) = 0$ we have $p_1(\Lambda_+^2 TM) = -c_2((\Lambda_+^2 TM)^c) = -c_2(\mathcal{C}\{\omega_J\} \oplus E_J^c) = c_1(E_J) \wedge c_1(E_J)$. QED

Remark 2. The canonical line bundle w.r.t J is the bundle $\mathcal{K}_J = \Lambda^2(TM^{(1,0)})^* = \Lambda^{(2,0)}$ of the complex 2-forms of type $(2, 0)$, and one has $(\Lambda_+^2 TM)^c = \mathcal{C}(\omega_J) \oplus \Lambda^{(2,0)} \oplus \Lambda^{(0,2)}$. The bundle E_J is isomorphic to the realification of the anti-canonical bundle $\mathcal{K}_J^{-1} = \Lambda^{(0,2)}$. As a complex line bundle, $\omega \rightarrow (\omega_c)^{(0,2)}$ is the complex isomorphism. Therefore, $c_1(E_J) = c_1(\mathcal{K}_J^{-1}) = -c_1(\mathcal{K}_J)$. An almost complex structure defines a canonic spin-c structure $\mathbf{s} = P_{Spin^c} M$ on M with canonic line bundle $\Lambda^{(2,0)}$ and complex spinor bundle $S = \Lambda^{(0,*)} = S^+ \oplus S^-$, $S^+ = \Lambda^{(0,0)} \oplus \Lambda^{(0,2)}$, $S^- = \Lambda^{(0,1)}$, where TM acts by Clifford multiplication. The Chern class of $\mathbf{s}$ is also given by $c_1(\mathcal{K}_J)$.

Finally we observe the following:

Lemma 3.1 *If J_1 and J_2 are two anti-commuting g -orthogonal almost Kähler complex structures on M , then $J_3 = J_1 J_2$ is also almost Kähler iff $g(\nabla_Z J_2(X), J_1 X) = 0 \ \forall X \in T_p M$. That is the case if either J_1 or J_2 is Kähler. In that case (J_1, J_2, J_3) is in fact an hyper-Kähler structure.*

Proof. It is sufficient to find an equivalent condition for $\delta\omega_3 = 0$. Using (3.1), (3.5), $\forall X \in T_p M$,

$$\begin{aligned} \nabla_{J_3 X} J_3(J_3 X) + \nabla_X J_3(X) &= \nabla_{J_1 J_2 X} (J_1 J_2)(J_3 X) + \nabla_X (J_1 J_2)(X) = \\ &= \nabla_{J_1 J_2 X} J_1(J_2 J_3 X) + J_1(\nabla_{J_1 J_2 X} J_2(J_3 X)) + \nabla_X J_1(J_2 X) + J_1(\nabla_X J_2(X)) \\ &= -J_1(\nabla_{J_2 X} J_1(J_2 J_3 X)) - J_1(\nabla_{J_2 J_1 X} J_2(J_3 X)) + \nabla_X J_1(J_2 X) + J_1(\nabla_X J_2(X)) \\ &= \nabla_{J_2 X} J_1(J_1 J_2 J_3 X) + J_1 J_2(\nabla_{J_1 X} J_2(J_3 X)) + \nabla_X J_1(J_2 X) + J_1(\nabla_X J_2(X)) \\ &= -\nabla_{J_2 X} J_1(X) + \nabla_X J_1(J_2 X) - J_1 \nabla_{J_1 X} J_2(J_2 J_3 X) + J_1(\nabla_X J_2(X)) \\ &= \left(-\nabla_{J_2 X} J_1(X) + \nabla_X J_1(J_2 X) \right) + J_1 \left(-\nabla_{J_1 X} J_2(J_1 X) + \nabla_X J_2(X) \right) \end{aligned}$$

Using (3.5) from $d\omega_1(X, J_2 X, Z) = 0 \ \forall X, Z$, we have $g(-\nabla_{J_2 X} J_1(X) + \nabla_X J_1(J_2 X), Z) = -g(\nabla_Z J_1(X), J_2 X)$. If X is a unit vector of $T_p M$ then $X, J_1 X, J_2 X, J_3 X$ is an o.n.b. Hence

$$\begin{aligned} -g(\delta J_3, Z) &= g(\nabla_X J_3(X) + \nabla_{J_3 X} J_3(J_3 X) + \nabla_{J_1 X} J_3(J_1 X) + \nabla_{J_3 J_1 X} J_3(J_3 J_1 X), Z) = \\ &= -g(\nabla_Z J_1(X), J_2 X) - g(-\nabla_{J_1 X} J_2(J_1 X) + \nabla_X J_2(X), J_1 Z) \\ &\quad -g(\nabla_Z J_1(J_1 X), J_2 J_1 X) - g(-\nabla_X J_2(X) + \nabla_{J_1 X} J_2(J_1 X), J_1 Z) \\ &= -g(\nabla_Z J_1(X), J_2 X) - g(J_1 \nabla_Z J_1(X), J_1 J_2 X) = -2g(\nabla_Z J_1(X), J_2 X). \end{aligned}$$

Assume that at a given point $\nabla X(p_0) = 0$. Thus at p_0 , $g(\nabla_Z J_1(X), J_2 X) = g(\nabla_Z(J_1(X)), J_2 X) = -g(J_1 X, \nabla_Z(J_2(X))) = -g(J_1 X, \nabla_Z J_2(X))$. Hitchin ([18], lemma (6.8)) proved that hyper-almost-Kähler structures are in fact hyper-Kähler. QED

4 The irreducible components of $R \in \mathcal{R}$

$\mathcal{R}$ has an orthogonal decomposition $\mathcal{R} = \oplus_{i=1}^4 \mathcal{R}_i$ into $O(V)$ -invariants subspaces ([25]), where

$$\begin{aligned}\mathcal{R}_1 &= (\text{Ker } b)^\perp = \{R : \sigma_R = 0\} = \{R : R = \lambda*\} \subset \text{Ker Ricci} \subset \text{Ker tr} \\ \mathcal{R}_2 &= (\text{Ker tr})^\perp = \{R = \lambda Id\} \\ \mathcal{R}_3 &= (\text{Ker } b) \cap (\text{Ker Ricci}) = \{R : *R = R*, \text{tr } R = \text{tr } *R = 0\} \\ \mathcal{R}_4 &= (\text{Ker } b) \cap (\text{Ker Ricci})^\perp = \{R : *R = -R*\}\end{aligned}$$

and they can be characterized in terms of the sectional and Ricci curvature. Set $\mathcal{B} := \text{Ker } b = \mathcal{R}_2 \oplus \mathcal{R}_3 \oplus \mathcal{R}_4$. We also have the following characterization using K_{isot} :

Proposition 4.1 *If $R \in \mathcal{R}$ and $b(R) = 0$ then $K_{isot} = 0$ iff $\sigma_R(P) = -\sigma_R(P^\perp) \forall P$. Hence,*

$$\mathcal{R}_4 = \{R : b(R) = 0, \sigma_R(P) = -\sigma_R(P^\perp) \forall P\} = \{R \in \mathcal{B} : K_{isot} = 0\} = \text{Sym}_0(V) \bullet g$$

and $\phi \rightarrow \phi \bullet g$ is an isomorphism from $\text{Sym}_0(V)$ onto $\mathcal{R}_4$.

Proof. In [25] it is proved the first equality of $\mathcal{R}_4$, and that the elements of this set satisfy $R(P, P^\perp) = 0$. Now it turns out that this condition and $\sigma_R(P) = -\sigma_R(P^\perp)$ is equivalent to $K_{isot} = 0$. We prove the less obvious implication. $K_{isot} = 0$ means that for any o.n.b. X, Y, Z, W , $2R(X, Y, Z, W) = R(X, Z, X, Z) + R(X, W, X, W) + R(Y, Z, Y, Z) + R(Y, W, Y, W)$. Setting $P = \text{span}\{X, Y\}$ and replacing X by $-X$ we conclude

$$\begin{aligned}0 &= R(X, Y, Z, W) = R(P, P^\perp) \\ 0 &= R(X, Z, X, Z) + R(X, W, X, W) + R(Y, Z, Y, Z) + R(Y, W, Y, W) =: \Psi(X, Y, Z, W).\end{aligned}$$

From $0 = \Psi(X, Y, Z, W) = \Psi(Z, Y, X, W) = \Psi(Z, X, W, Y)$ we have $0 = \Psi(Z, Y, X, W) - \Psi(X, Y, Z, W) + \Psi(Z, X, W, Y) = 2R(Z, W, Z, W) + 2R(X, Y, X, Y) = 2\sigma_R(P^\perp) + 2\sigma_R(P)$. For a proof that $\phi \rightarrow \phi \bullet g$ defines an isomorphism see e.g. [25]. QED

Let $\phi \in \text{Sym}(V)$, and e_i a d.o.n.b. of eigenvectors of ϕ , with corresponding eigenvalues λ_i . Then $e_i \wedge e_j$ with $i < j$ are the eigenvectors of $\phi \bullet g$ ($\phi \bullet \phi$ resp.) corresponding to the respective eigenvalues $\lambda_i + \lambda_j$ ($2\lambda_i \lambda_j$ resp.). Let $\Lambda_1^\pm = e_1 \wedge e_2 \pm e_3 \wedge e_4$, $\Lambda_2^\pm = e_1 \wedge e_3 \mp e_2 \wedge e_4$, $\Lambda_3^\pm = e_1 \wedge e_4 \pm e_2 \wedge e_3$. We have:

$$\begin{aligned}\langle \phi \bullet g(\Lambda_\alpha^\pm), \Lambda_\beta^\pm \rangle &= \delta_{\alpha\beta} \text{tr}(\phi) & \langle \phi \bullet g(\Lambda_1^+), \Lambda_\beta^- \rangle &= \delta_{1\beta}(\lambda_1 + \lambda_2 - \lambda_3 - \lambda_4) \\ \langle \phi \bullet g(\Lambda_2^+), \Lambda_\beta^- \rangle &= \delta_{2\beta}(\lambda_1 + \lambda_3 - \lambda_2 - \lambda_4) & \langle \phi \bullet g(\Lambda_3^+), \Lambda_\beta^- \rangle &= \delta_{3\beta}(\lambda_1 + \lambda_4 - \lambda_2 - \lambda_3) \\ \langle \phi \bullet \phi(\Lambda_1^\pm), \Lambda_\beta^\pm \rangle &= 2\delta_{1\beta}(\lambda_1 \lambda_2 + \lambda_3 \lambda_4) & \langle \phi \bullet \phi(\Lambda_1^\pm), \Lambda_\beta^\mp \rangle &= 2\delta_{1\beta}(\lambda_1 \lambda_2 - \lambda_3 \lambda_4) \\ \langle \phi \bullet \phi(\Lambda_2^\pm), \Lambda_\beta^\pm \rangle &= 2\delta_{2\beta}(\lambda_1 \lambda_3 + \lambda_2 \lambda_4) & \langle \phi \bullet \phi(\Lambda_2^\pm), \Lambda_\beta^\mp \rangle &= 2\delta_{2\beta}(\lambda_1 \lambda_3 - \lambda_2 \lambda_4) \\ \langle \phi \bullet \phi(\Lambda_3^\pm), \Lambda_\beta^\pm \rangle &= 2\delta_{3\beta}(\lambda_1 \lambda_4 + \lambda_2 \lambda_3) & \langle \phi \bullet \phi(\Lambda_3^\pm), \Lambda_\beta^\mp \rangle &= 2\delta_{3\beta}(\lambda_1 \lambda_4 - \lambda_2 \lambda_3).\end{aligned}$$

Lemma 4.1 $(\phi \bullet g)_{\pm\pm}^\pm = \frac{1}{2} \text{tr}(\phi) Id_{\Lambda_2^\pm}$, and $\text{tr}(\phi \bullet \phi) = \sum_{i < j} 2\lambda_i \lambda_j = 2\sigma_2(\phi)$.

Lemma 4.2 Let $\phi, \xi \in \text{Sym}(V)$ and $\langle \phi, \xi \rangle = \sum_{ij} \phi(e_i, e_j) \xi(e_i, e_j)$ where e_i is an o.n.b. of V . Then

$$\begin{aligned}\xi \bullet \phi &\in \mathcal{B} & \text{tr}(\phi \bullet g) &= \langle Id, \phi \bullet g \rangle = 3\text{tr}(\phi) \\ \langle \phi \bullet g, \xi \bullet g \rangle &= 2\langle \phi, \xi \rangle + \text{tr}(\phi) \text{tr}(\xi) & *(\phi \bullet g) &= \frac{1}{2} \text{tr}(\phi) g \bullet g - \phi \bullet g \\ \text{Ricci}_{\phi \bullet g} &= \text{Tr}(\phi) g + 2\phi & s_{\phi \bullet g} &= 6\text{tr} \phi \\ \langle \phi \bullet g, * \xi \bullet g \rangle &= 0 & \phi \bullet g &= \xi \bullet g \text{ iff } \phi = \xi.\end{aligned}$$

Proof. The equalities are proved using the eigenvalues λ_i and eigenvectors e_i of ϕ . Since both $P = e_i \wedge e_j$ and $P^\perp$ are eigenvectors of $R = \phi \bullet g$, then $*R*$ has the same eigenvalues $\lambda'_i + \lambda'_j$ as R . Namely, if $(ijkl)$ is a permutation of 1234, $\lambda'_i + \lambda'_j = \sum_s \lambda_s - (\lambda_i + \lambda_j) = \text{tr}(\phi) - (\lambda_i + \lambda_j) = \lambda_k + \lambda_l$. QED

If $R \in \mathcal{R}$, then $R_1 = \frac{1}{3}t(R)* = \frac{1}{6}\text{tr}(*R)*$. For $R \in \mathcal{B}$ we have the decomposition:

$$R_2 = \frac{1}{12}\text{tr}(R)g \bullet g = \frac{1}{24}s_R g \bullet g = \frac{1}{12}s_R Id \quad (4.1)$$

$$R_3 = \frac{1}{2}(R + *R*) - \frac{1}{12}\text{tr}(R)g \bullet g = \mathcal{W} = \mathcal{W}^+ + \mathcal{W}^- \quad (4.2)$$

$$R_4 = \frac{1}{2}(R - *R*) = \frac{1}{2}(\text{Ricci}_R - \frac{1}{4}s_R g) \bullet g. \quad (4.3)$$

$\mathcal{W}$ is the Weyl tensor. It applies $\bigwedge_\pm^2 V$ into $\bigwedge_\pm^2 V$, with $\mathcal{W}^+$ and $\mathcal{W}^-$ the self-dual and the anti-selfdual part respectively, i.e, they satisfy: $*\mathcal{W}^\pm = \mathcal{W}^\pm* = \pm\mathcal{W}^\pm$. Furthermore $R_\pm^\pm = \mathcal{W}^\pm + \frac{s_R}{12}Id_{\bigwedge_\pm^2}$, $R_-^+ + R_+^- = \frac{1}{2}(\text{Ricci} - \frac{s_R}{4}g) \bullet g$. From previous lemmas we have:

Proposition 4.2 $\mathcal{R}_4 \oplus \mathcal{R}_2 = \text{Sym}(V) \bullet g$, and $\forall \phi \in \text{Sym}(V)$ $(\phi \bullet g)_2 = \frac{\text{tr}(\phi)}{4}g \bullet g$, $(\phi \bullet g)_4 = (\phi - \frac{\text{tr}(\phi)}{4}g) \bullet g$.

Lemma 4.3 Let $R \in \mathcal{B}$ and $\phi, \xi \in \text{Sym}(V)$. Then $\langle R, \phi \bullet g \rangle = \langle \text{Ricci}_R, \phi \rangle$, $\langle *R*, \phi \bullet g \rangle = \frac{1}{2}s_R \text{tr}(\phi) - \langle \text{Ricci}_R, \phi \rangle$, and $\langle *\phi \bullet g*, \phi \bullet g \rangle = 2(\text{tr}(\phi))^2 - 2\|\phi\|^2 = 2\text{tr}(\phi \bullet \phi)$.

Proof. Using the decomposition (4.1)-(4.3), lemma 4.2 and proposition 4.2 we have

$$\begin{aligned} \langle R, \phi \bullet g \rangle &= \frac{1}{24}s_R \langle g \bullet g, \phi \bullet g \rangle + \frac{1}{2} \langle (\text{Ricci}_R - \frac{1}{4}s_R g) \bullet g, \phi \bullet g \rangle \\ &= \frac{1}{24}s_R (2\langle g, \phi \rangle + \text{tr}(g)\text{tr}(\phi)) + \langle (\text{Ricci}_R - \frac{1}{4}s_R g), \phi \rangle = \langle \text{Ricci}_R, \phi \rangle. \end{aligned}$$

$$\langle *R*, \phi \bullet g \rangle = \langle \text{Ricci}_{*R*}, \phi \rangle = \langle \frac{1}{2}s_R g - \text{Ricci}_R, \phi \rangle = \frac{1}{2}s_R \text{tr}(\phi) - \langle \text{Ricci}_R, \phi \rangle. \quad \text{QED.}$$

Proposition 4.3 If $Q = R + \phi \bullet g$, where $R \in \mathcal{B}$ and $\phi \in \text{Sym}(V)$, then

$$4\pi^2 \mathcal{X}(Q) = 4\pi^2 \mathcal{X}(R) + \left(\frac{s_R}{2} \text{tr}(\phi) - \langle \text{Ricci}_R, \phi \rangle + (\text{tr}(\phi))^2 - \|\phi\|^2 \right) \text{Vol}_V \quad (4.4)$$

$$4\pi^2 p_1(Q) = 4\pi^2 p_1(R). \quad (4.5)$$

Proof. From lemma 4.3

$$\begin{aligned} 4\pi^2 \langle \mathcal{X}(Q), \text{Vol}_V \rangle &= \frac{1}{2} \langle *Q*, Q \rangle = \frac{1}{2} \langle *R*, R \rangle + \langle *R*, \phi \bullet g \rangle + \frac{1}{2} \langle *(\phi \bullet g)*, \phi \bullet g \rangle \\ &= 4\pi^2 \langle \mathcal{X}(R), \text{Vol}_V \rangle + \frac{s_R}{2} \text{tr}(\phi) - \langle \text{Ricci}_R, \phi \rangle + (\text{tr}(\phi))^2 - \|\phi\|^2. \end{aligned}$$

Note that $R = R^T$, $Q = Q^T$. Using decomposition (4.1)(4.2)(4.3) and lemmas 4.2, 4.1, we have

$$\begin{aligned} 4\pi^2 \langle p_1(Q), \text{Vol}_V \rangle &= \langle Q, *Q \rangle = \langle R, *R \rangle + 2\langle R, *\phi \bullet g \rangle + \langle \phi \bullet g, *\phi \bullet g \rangle \\ &= 4\pi^2 \langle p_1(R), \text{Vol}_V \rangle + 2\langle \mathcal{W}, *\phi \bullet g \rangle = 4\pi^2 p_1(R) + 2\langle *\mathcal{W}, \phi \bullet g \rangle \\ &= 4\pi^2 \langle p_1(R), \text{Vol}_V \rangle + 2\langle \mathcal{W}^+, \phi \bullet g \rangle - 2\langle \mathcal{W}^-, \phi \bullet g \rangle \\ &= 4\pi^2 \langle p_1(R), \text{Vol}_V \rangle + 2\langle \mathcal{W}^+, (\phi \bullet g)_+^\perp \rangle - 2\langle \mathcal{W}^-, (\phi \bullet g)_-^\perp \rangle \\ &= 4\pi^2 \langle p_1(R), \text{Vol}_V \rangle + \text{tr}(\phi) (\langle \mathcal{W}^+, Id_{\bigwedge_+^2} \rangle - \langle \mathcal{W}^-, Id_{\bigwedge_-^2} \rangle) \\ &= 4\pi^2 \langle p_1(R), \text{Vol}_V \rangle + \text{tr}(\phi) (\text{tr}(\mathcal{W}^+) - \text{tr}(\mathcal{W}^-)). \end{aligned}$$

Since $\mathcal{W}$ and $*\mathcal{W}$ have zero trace, the same holds for $\mathcal{W}^+$ and $\mathcal{W}^-$ leading to (4.5). QED

Corollary 4.1 *If λ_i are the eigenvalues of ϕ then $\mathcal{X}(\phi \bullet g) = 0$ iff $\sigma_2(\phi) := \sum_{i < j} \lambda_i \lambda_j = 0$.*

Proposition 4.4 *If $R \in \mathcal{B}$ and $\lambda \in \mathbf{R}$, then*

$$\begin{aligned} 4\pi^2 \mathcal{X}(R + \lambda*) &= 4\pi^2 \mathcal{X}(R) + 3\lambda^2 \text{Vol}_V \\ 4\pi^2 p_1(R + \lambda*) &= 4\pi^2 p_1(R) + \lambda s_R \text{Vol}_V. \end{aligned}$$

Remark 3. Using (4.1)-(4.3) one sees that for $R \in \mathcal{B}$ (see e.g. [7])

$$\begin{aligned} 8\pi^2 \mathcal{X}(R) &= (\|R_2\|^2 + \|R_3\|^2 - \|R_4\|^2) \text{Vol}_V \\ 4\pi^2 p_1(R) &= \langle R_3, R_3* \rangle \text{Vol}_V = (\|\mathcal{W}^+\|^2 - \|\mathcal{W}^-\|^2) \text{Vol}_V. \end{aligned}$$

The invariance of the Pontrjagin class under a conformal change of the metric now follows from the well known conformal invariance of the Weyl tensor, but we give a simple proof in Prop.4.3.

5 Conformal change of the metric

Let g be a Riemannian metric on a 4-dimensional manifold M and $f : M \rightarrow \mathbf{R}$ a smooth map. We denote by ∇ and $\hat{\nabla}$ the Levi Civita connections of g and $\hat{g} = e^f g$ with curvature tensors $R^M, \hat{R}^M$ respectively. Then $\hat{\nabla}_X Y = \nabla_X Y + \hat{S}(X, Y)$ where $\hat{S}(X, Y) = \frac{1}{2}(df(X)Y + df(Y)X - \langle X, Y \rangle \nabla f)$. The curvature tensors $\hat{R}^M(X, Y, Z, W) = \hat{g}(\hat{R}^M(X, Y)Z, W)$, and $R^M(X, Y, Z, W) = g(R^M(X, Y)Z, W)$ are related by

$$\hat{R}^M(X, Y, Z, W) = e^f R^M(X, Y, Z, W) + e^f \phi \bullet g(X, Y, Z, W) \quad (5.1)$$

$$\text{where } \phi = \frac{1}{2} \left(-\frac{1}{4} \|\nabla f\|^2 g + \frac{1}{2} df \otimes df - \text{Hess}(f) \right). \quad (5.2)$$

Let $\text{Hess}(f)(X, Y) = \nabla_X df(Y) = \nabla_Y df(X)$ be the Hessian of f , $\Delta df = (d\delta + \delta d)df = d\delta df$, and $S(df)(Y) = df(\text{Ricci}^M(Y)) = \text{Ricci}^M(Y, \nabla f)$, the Weitzenböck operator S on the 1-form df . We are using the following sign for the Laplacian of maps $h : M \rightarrow \mathbf{R}$, $\Delta h = \Delta^+ h = -\delta dh = \text{div}(\nabla h) = \text{tr}(\text{Hess}(h))$. Set $\nabla_{X,Y}^2 df = \nabla_X(\nabla_Y df) - \nabla_{\nabla_X Y} df$.

Lemma 5.1 *For all $X, Y, Z \in T_p M$, $p \in M$*

- (a) $\nabla_{X,Y}^2 df(Z) = \nabla_{X,Z}^2 df(Y)$.
- (b) $\nabla_{X,Y}^2 df(Z) - \nabla_{Y,X}^2 df(Z) = df(R^M(X, Y)Z) = \nabla_{X,Z}^2 df(Y) - \nabla_{Y,Z}^2 df(X)$.
- (c) $\Delta df = -d(\Delta f)$.
- (d) $-\frac{1}{2} \Delta \|\nabla f\|^2 = \langle \Delta df, df \rangle - \|\text{Hess}(f)\|^2 - \text{Ricci}^M(\nabla f, \nabla f)$.
- (e) $(\Delta f)^2 = -\frac{1}{2} \Delta \|\nabla f\|^2 + \text{div}(\Delta f \nabla f) + \|\text{Hess}(f)\|^2 + \text{Ricci}^M(\nabla f, \nabla f)$.
- (f) $\|\nabla f\|^2 \Delta f = \text{div}(\|\nabla f\|^2 \nabla f) - 2\text{Hess}(f)(\nabla f, \nabla f)$.

Proof. We may assume without loss of generality that $\nabla X = \nabla Y = \nabla Z = 0$ at a given p_0 . From $\nabla_X(\nabla_Z df(Y)) = \nabla_X(\nabla_Y df(Z))$, we get (a) at p_0 . At p_0

$$\begin{aligned}
\nabla_X (\nabla_Y df(Z)) - \nabla_Y (\nabla_X df(Z)) &= \\
&= \nabla_X (\nabla_Y (df(Z))) - \nabla_Y (\nabla_X (df(Z))) - \nabla_X (df(\nabla_Y Z)) + \nabla_Y (df(\nabla_X Z)) \\
&= df(-\nabla_X \nabla_Y Z + \nabla_Y \nabla_X Z) = df(R^M(X, Y)Z).
\end{aligned}$$

Thus (b) is proved. By (b), if $e_1, \dots, e_4$ is any o.n. frame of M with $\nabla e_i(p_0) = 0$, then at p_0

$$df(Ricci^M(Y)) = \sum_i \nabla_{e_i, e_i}^2 df(Y) - \nabla_{Y, e_i}^2 df(e_i) = \sum_i \nabla_{e_i, e_i}^2 df(Y) - \nabla_Y(\Delta f).$$

Consequently, by the Weitzenböck formula

$$\Delta df(Y) = (d\delta + \delta d)df(Y) = -(Trace \nabla^2 df)(Y) + S(df)(Y) = -\nabla_Y(\Delta f).$$

So, we have proved (c). (d) is a well-known application of the Weitzenböck formula for 1-forms. (e) and (f) follows from previous alignments and that $div(hX) = \langle X, \nabla h \rangle + h div(X)$. QED

Lemma 5.2

$$tr(\phi) = -\frac{1}{4}\|\nabla f\|^2 - \frac{1}{2}\Delta f \quad (5.3)$$

$$tr(\phi \bullet \phi) = (tr(\phi))^2 - \|\phi\|^2 = \frac{1}{8}div((2\Delta f + \|\nabla f\|^2)\nabla f - \nabla(\|\nabla f\|^2)) + \frac{1}{4}Ricci^M(\nabla f, \nabla f) \quad (5.4)$$

$$\frac{1}{2}s^M tr(\phi) - \langle Ricci^M, \phi \rangle = -\frac{1}{4}s^M \Delta f - \frac{1}{4}Ricci^M(\nabla f, \nabla f) + \frac{1}{2}\langle Ricci^M, Hess(f) \rangle. \quad (5.5)$$

Proof. Let e_i be an o.n.b. of $T_p M$. We have

$$\begin{aligned}
tr(\phi) &= \frac{1}{2}(-\|\nabla f\|^2 + \sum_i \frac{1}{2}df(e_i)df(e_i) - Hess(f)(e_i, e_i)) = -\frac{1}{4}\|\nabla f\|^2 - \frac{1}{2}\Delta f \\
\|\phi\|^2 &= \frac{1}{4}\|Hess f\|^2 + \frac{1}{8}\|\nabla f\|^2 \Delta f - \frac{1}{4}Hess(f)(\nabla f, \nabla f) + \frac{1}{16}\|\nabla f\|^4.
\end{aligned}$$

Thus

$$(tr(\phi))^2 - \|\phi\|^2 = \frac{1}{4}(\Delta f)^2 + \frac{1}{8}\|\nabla f\|^2 \Delta f - \frac{1}{4}\|Hess(f)\|^2 + \frac{1}{4}Hess(f)(\nabla f, \nabla f).$$

Using Lemma 5.1 (e) and (f) we get (5.4). Now

$$\begin{aligned}
\frac{1}{2}s^M tr(\phi) - \langle Ricci^M, \phi \rangle &= \\
&= \frac{1}{2}s^M(-\frac{1}{4}\|\nabla f\|^2 - \frac{1}{2}\Delta f) + \frac{1}{8}\|\nabla f\|^2 s^M - \frac{1}{4}Ricci^M(\nabla f, \nabla f) + \frac{1}{2}\langle Ricci^M, Hess(f) \rangle \\
&= -\frac{1}{4}s^M \Delta f - \frac{1}{4}Ricci^M(\nabla f, \nabla f) + \frac{1}{2}\langle Ricci^M, Hess(f) \rangle. \quad \text{QED}
\end{aligned}$$

We recall the well-known formula (see e.g. [12]) $\delta(Ricci^M)^\sharp + \frac{1}{2}\nabla s^M = 0$, derived from the second Bianchi identity, where δ is the formal adjoint to d for TM -valued forms on M . Now

$$\begin{aligned}
div((Ricci^M)^\sharp(\nabla f)) &= \sum_i \langle \nabla_{e_i}((Ricci^M)^\sharp(\nabla f)), e_i \rangle \\
&= \sum_i \langle (\nabla_{e_i}(Ricci^M)^\sharp)(\nabla f), e_i \rangle + \langle (Ricci^M)^\sharp(\nabla_{e_i} \nabla f), e_i \rangle \\
&= -\langle \delta((Ricci^M)^\sharp), \nabla f \rangle + \langle Ricci^M, Hess(f) \rangle.
\end{aligned}$$

Moreover, $g(\nabla s^M, \nabla f) = \langle ds^M, df \rangle = d(s^M * df) + s^M \delta df = \text{div}(s^M \nabla f) - s^M \Delta f$. Thus

$$\begin{aligned} 0 &= g(\delta(\text{Ricci}^M)^\sharp + \tfrac{1}{2} \nabla s^M, \nabla f) \\ &= \text{div}(-(\text{Ricci}^M)^\sharp(\nabla f) + \tfrac{1}{2} s^M \nabla f) + \langle \text{Ricci}^M, \text{Hess}(f) \rangle - \tfrac{1}{2} s^M \Delta f. \end{aligned}$$

So we have obtained:

Lemma 5.3 $\langle \text{Ricci}^M, \text{Hess}(f) \rangle - \tfrac{1}{2} s^M \Delta f = \text{div}(-\tfrac{1}{2} s^M \nabla f + (\text{Ricci}^M)^\sharp(\nabla f)).$

Proof of Proposition 1.1. For each vector field Z on M we denote $\hat{Z} = e^{-\frac{f}{2}} Z$. The curvature tensor $\hat{R}^M$ of TM w.r.t the Levi-Civita connection of the Riemannian metric $\hat{g}$ can be seen as a bundle map $Q : \wedge^2 TM \rightarrow \wedge^2 TM$ given by $g(Q(X \wedge Y), (Z \wedge W)) = Q(X, Y, Z, W) = \hat{R}^M(X, Y, \hat{Z}, \hat{W})$. Thus $Q(X, Y, Z, W) = R^M(X, Y, Z, W) + \phi \bullet g(X, Y, Z, W)$, where ϕ is given by (5.2). Therefore $\mathcal{X}(\hat{R}^M) = \mathcal{X}(Q) = \mathcal{X}(R^M + \phi \bullet g)$. Now we apply Proposition 4.3 and Lemmas 5.2 and 5.3 to obtain (1.2). Similar for $p_1(\hat{R}^M)$. QED

6 Connections with torsion

Let (M, g) be a manifold of dimension 4 and ∇ be its Levi-Civita connection, and ∇' any other connection. Set $T(X, Y) = \nabla'_X Y - \nabla'_Y X - [X, Y]$, and $S(X, Y) = \nabla'_X Y - \nabla_X Y$. Then $T(X, Y) = S(X, Y) - S(Y, X)$. From $\nabla'_X g(Y, Z) = g(S(X, Y), Z) + g(S(X, Z), Y)$, we see that ∇' is a g -Riemannian connection iff $g(S(X, Y), Z) = -g(S(X, Z), Y)$. Moreover $\nabla' = \nabla$ iff $T = 0$, that is $S(X, Y) = S(Y, X)$. If $\zeta : TM \rightarrow E$ is a vector bundle map and $E \rightarrow M$ has a connection ∇^E then $d\zeta = d\nabla^E \zeta$ is a 2-form on M with values on E , $d\zeta(X, Y) = \nabla_X^E(\zeta(Y)) - \nabla_Y^E(\zeta(X)) - \zeta([X, Y])$. Since ∇ is a torsion free connection on M , $d\zeta(X, Y) = \nabla_X \zeta(Y) - \nabla_Y \zeta(X)$. We define $\mathcal{S} : TM \rightarrow L(TM; TM)$ and $\mathcal{S}^2 : \wedge^2 TM \rightarrow L(TM; TM)$ by

$$\mathcal{S}(X)(Y) = S(X, Y) \quad \mathcal{S}^2(X \wedge Y)(Z) = S(X, S(Y, Z)) - S(Y, S(X, Z))$$

and take the exterior derivative $d\mathcal{S} : \wedge^2 TM \rightarrow L(TM; TM)$, by taking the usual connection on $L(TM; TM)$ w.r.t the Levi-Civita connection of M . We have

$$R'(X \wedge Y)(Z) = R^M(X \wedge Y)(Z) - d\mathcal{S}(X \wedge Y)(Z) - \mathcal{S}^2(X \wedge Y)(Z).$$

If ∇' is g -Riemannian then $\mathcal{S}$, $d\mathcal{S}$ and $\mathcal{S}^2$ take values on $\wedge^2 TM$ and $\langle \mathcal{S}(X), Y \wedge Z \rangle = g(S(X, Y), Z)$,

$$\begin{aligned} \langle \mathcal{S}^2(X \wedge Y), Z \wedge W \rangle &= g(S(X, Z), S(Y, W)) - g(S(X, W), S(Y, Z)) \\ \langle R'(X \wedge Y), Z \wedge W \rangle &= \langle R^M(X \wedge Y), Z \wedge W \rangle - \langle d\mathcal{S}(X \wedge Y), Z \wedge W \rangle - \langle \mathcal{S}^2(X \wedge Y), Z \wedge W \rangle. \end{aligned}$$

7 Transgression forms

Let (M, g) be a Riemannian manifold of dimension 4 with Levi-Civita connection ∇ and let ∇' be any other g -Riemannian connection. We take $\pi : \tilde{M} = M \times [0, 1] \rightarrow M$ the projection map $\pi(p, t) = p$ and the connection $\tilde{\nabla}$ on $\pi^{-1}TM$ given by $\tilde{\nabla} = t\nabla + (1-t)\nabla'$, where ∇ and ∇' are the

pullback connections w.r.t π . For any vector fields X, Y, Z, W of M , map $f : M \times [0, 1] \rightarrow \mathbf{R}$ and $u \in T_p M$, $h \in \mathbf{R}$, $\tilde{\nabla}_{(u,h)}(f\pi^{-1}Y)_{(p,t)} = df(p,t)(u,h)Y_p + f(p,t)\nabla_u Y_{(p)} + tf(p,t)S_p(u, Y_p)$. Thus

$$\tilde{\nabla}_{\frac{d}{dt}}\tilde{\nabla}_{(X,0)}(\pi^{-1}Y) = \tilde{\nabla}_{\frac{d}{dt}}(\pi^{-1}(\nabla_X Y) + t\pi^{-1}(S(X, Y))) = S(X, Y)$$

and assuming at a given point p , $\nabla Y_p = \nabla Z_p = 0$, then at (p, t)

$$\tilde{\nabla}_{(X,0)}\tilde{\nabla}_{(Y,0)}(\pi^{-1}Z) = \nabla_X \nabla_Y Z + t\nabla_X S(Y, Z) + t^2 S(X, S(Y, Z)).$$

Therefore, the curvature tensor of $\pi^{-1}TM$ with respect to this connection (Riemannian w.r.t $\pi^{-1}g$), $\tilde{R} : \wedge^2 \pi^{-1}TM \rightarrow \wedge^2 T\tilde{M}$ $\tilde{R}(\pi^{-1}Z, \pi^{-1}W)(\tilde{X}, \tilde{Y}) =: \tilde{R}(\tilde{X}, \tilde{Y}, \pi^{-1}Z, \pi^{-1}W)$ satisfies

$$\begin{aligned} \tilde{R}(X, Y, Z, W) &= R^M(X, Y, Z, W) - tdS(X \wedge Y)(Z \wedge W) - t^2 S^2(X \wedge Y)(Z \wedge W) \\ \tilde{R}(\frac{d}{dt}, Y, Z, W) &= -S(Y)(Z \wedge W) \end{aligned}$$

with $dS = d^\nabla S$ where ∇ denotes de induced connection on $\wedge^2 TM$ by ∇ . Let e_1, e_2, e_3, e_4 be an d.o.n.b. of $T_p M = (\pi^{-1}TM)_{(p,t)}$. The definitions of $\mathcal{X}(\tilde{R})$ and $p_1(\tilde{R})$ are the 4-forms on $\tilde{M}$

$$\mathcal{X}(\tilde{R}) = \sum_{i < j} \frac{1}{8\pi^2} \tilde{R}(e_i \wedge e_j) \wedge \tilde{R}(*e_i \wedge e_j) \quad p_1(\tilde{R}) = \sum_{i < j} \frac{1}{4\pi^2} \tilde{R}(e_i \wedge e_j) \wedge \tilde{R}(e_i \wedge e_j).$$

For $\zeta : V \rightarrow \wedge^2 V$ and $R \in L(\wedge^2 V; \wedge^2 V)$, we define a 3-form

$$\langle \zeta \wedge R \rangle(X, Y, Z) := \langle \zeta(X), R(Y, Z) \rangle + \langle \zeta(Z), R(X, Y) \rangle + \langle \zeta(Y), R(Z, X) \rangle. \quad (7.1)$$

Now we compute $\mathcal{X}(\tilde{R})(\frac{d}{dt}, X, Y, Z)$.

$$\begin{aligned} &8\pi^2 \mathcal{X}(\tilde{R})(\frac{d}{dt}, X, Y, Z) = \\ &= \sum_{i < j} \left(\langle \tilde{R}(\frac{d}{dt}, X), e_i \wedge e_j \rangle \langle \tilde{R}(Y, Z), *e_i \wedge e_j \rangle - \langle \tilde{R}(\frac{d}{dt}, Y), e_i \wedge e_j \rangle \langle \tilde{R}(X, Z), *e_i \wedge e_j \rangle \right. \\ &\quad + \langle \tilde{R}(\frac{d}{dt}, Z), e_i \wedge e_j \rangle \langle \tilde{R}(X, Y), *e_i \wedge e_j \rangle + \langle \tilde{R}(Y, Z), e_i \wedge e_j \rangle \langle \tilde{R}(\frac{d}{dt}, X), *e_i \wedge e_j \rangle \\ &\quad \left. - \langle \tilde{R}(X, Z), e_i \wedge e_j \rangle \langle \tilde{R}(\frac{d}{dt}, Y), *e_i \wedge e_j \rangle + \langle \tilde{R}(X, Y), e_i \wedge e_j \rangle \langle \tilde{R}(\frac{d}{dt}, Z), *e_i \wedge e_j \rangle \right) \\ &= \sum_{i < j} 2 \left(-S(X)(e_i \wedge e_j) \left(\langle R^M(Y, Z), *e_i \wedge e_j \rangle - t \langle dS(Y \wedge Z), *e_i \wedge e_j \rangle - t^2 \langle S^2(Y \wedge Z), *e_i \wedge e_j \rangle \right) \right. \\ &\quad + S(Y)(e_i \wedge e_j) \left(\langle R^M(X, Z), *e_i \wedge e_j \rangle - t \langle dS(X \wedge Z), *e_i \wedge e_j \rangle - t^2 \langle S^2(X \wedge Z), *e_i \wedge e_j \rangle \right) \\ &\quad \left. - S(Z)(e_i \wedge e_j) \left(\langle R^M(X, Y), *e_i \wedge e_j \rangle - t \langle dS(X \wedge Y), *e_i \wedge e_j \rangle - t^2 \langle S^2(X \wedge Y), *e_i \wedge e_j \rangle \right) \right) \\ &= -2 \langle S \wedge * (R^M - tdS - t^2 S^2) \rangle(X, Y, Z). \end{aligned}$$

Similarly, $4\pi^2 p_1(\tilde{R})(\frac{d}{dt}, X, Y, Z) = -2 \langle S \wedge (R^M - tdS - t^2 S^2) \rangle(X, Y, Z)$. Integration over $[0, 1]$ gives:

Proposition 7.1 *If (M, g) is an oriented Riemannian manifold of dimension 4 with Levi-Civita connection ∇ , and $\nabla_X Y = \nabla_X Y + S(X, Y)$ is a g -Riemannian connection on TM with torsion $T(X, Y) = S(X, Y) - S(Y, X)$ and curvature tensor R' , then*

$$\mathcal{X}(R') = \mathcal{X}(R^M) - \frac{1}{4\pi^2} d \left(\langle S \wedge * (R^M - \frac{1}{2} dS - \frac{1}{3} S^2) \rangle \right) \quad (7.2)$$

$$p_1(R') = p_1(R^M) - \frac{1}{2\pi^2} d \left(\langle S \wedge (R^M - \frac{1}{2} dS - \frac{1}{3} S^2) \rangle \right). \quad (7.3)$$

Remark 4. The Chern-Simons transgression formula is a particular case of the following fact: If φ is a closed k -form on $M \times [a, b]$ and $\varphi_t = i_t^* \varphi$, where $i_t : M \rightarrow M \times [a, b]$, $i_t(p) = (p, t)$, then $\varphi_b - \varphi_a = dT$ where T is the $(k-1)$ -form on M given by: $T(X_1, \dots, X_{k-1}) = \int_a^b \varphi(\frac{d}{dt}, (X_1, 0), \dots, (X_{k-1}, 0)) dt$.

8 Singular connections

Let (M, g) be a Riemannian 4-manifold with Levi-Civita connection ∇ and (E, g_E, ∇^E) a Riemannian vector bundle of rank 4 and Riemannian connection ∇^E and assume there exist a vector bundle map $\Phi : TM \rightarrow E$ that is conformal with $g_E(\Phi(X), \Phi(Y)) = h(p)g(X, Y) \forall X, Y \in T_p M$, where $h : M \rightarrow \mathbf{R}$ is a smooth non-negative function with zero set Σ . On TM we have a degenerated metric $\hat{g} = hg$, with Levi-Civita connection $\hat{\nabla}$, defined on $M \sim \Sigma$ and let ∇' be the $\hat{g}$ -Riemannian connection on TM , defined along $M \sim \Sigma$, and that makes $\Phi : TM \rightarrow E$ a parallel isometry. We define $\hat{S}(X, Y) = \hat{\nabla}_X Y - \nabla_X Y$, $S'(X, Y) = \nabla'_X Y - \hat{\nabla}_X Y$, and $S(X, Y) = \nabla'_X Y - \nabla_X Y = S'(X, Y) + \hat{S}(X, Y)$. The connection ∇' satisfies $\Phi(\nabla'_X Y) = \nabla_X^E(\Phi(Y))$, and may not be smoothly extended to Σ , defining a singular connection. But the curvature tensor $R' : \wedge^2 TM \rightarrow \wedge^2 TM$ satisfies

$$\begin{aligned} R'(X, Y, Z, W) &= \hat{g}(R'(X, Y)Z, W) = hg(R'(X, Y)Z, W) = g_E(\Phi(R'(X, Y)Z), \Phi(W)) \\ &= g_E(R^E(X, Y)(\Phi(Z)), \Phi(W)) = R^E(X, Y, \Phi(Z), \Phi(W)) \end{aligned}$$

where $R^E : \wedge^2 TM \rightarrow \wedge^2 E$ is the curvature tensor of (E, g_E, ∇^E) . The above equality means that we can smoothly extend R' as a 4-tensor to all M , defining it to be zero along Σ . Moreover, we can define $\mathcal{X}(R')$ as $\mathcal{X}(R^E)$ on all M , and similar for $p_1(R')$. Note that the star operator $* : \wedge^2 TM \rightarrow \wedge^2 TM$ is the same for any conformal change of the metric. Now we have $\Phi(S(X, Y)) = \Phi(\nabla'_X Y) - \Phi(\nabla_X Y) = \nabla_X^E(\Phi(Y)) - \Phi(\nabla_X Y) = \nabla_X \Phi(Y)$. Therefore

$$S = \Phi^{-1} \circ \nabla \Phi \quad T = \Phi^{-1} \circ d\Phi \quad (8.1)$$

$$S' = S - \hat{S} = \Phi^{-1} \circ \nabla \Phi - \frac{1}{2} d \log h \odot Id + \frac{1}{2} g \otimes \nabla \log h \quad (8.2)$$

and the torsion $T' = T$. From the two previous sections we obtain the formulae in Theorem 1.1.

Proposition 8.1 *In the condition of Theorem 1.1, if $d\Phi = 0$, then $\mathcal{X}(R^E) = \mathcal{X}(R^M) + \frac{1}{32\pi^2} \text{div}_g(P(\nabla \log h)) \text{Vol}_M$ and $p_1(R^E) = p_1(R^M)$. Moreover, in this case, $\delta\Phi = 0$ iff h is constant, and so $\mathcal{X}(R^E) = \mathcal{X}(R^M)$. This is the case when Φ is harmonic and M is compact.*

Proof. If $d\Phi = 0$, from (8.1), $T' = T = 0$, and so $\nabla' = \hat{\nabla}$. Hence, $S' = 0$, that is $\Phi^{-1} \nabla_X \Phi(Y) = \frac{1}{2} d \log h(X)Y + \frac{1}{2} d \log h(Y)X - \frac{1}{2} g(X, Y) \nabla \log h$. Thus, $\Phi^{-1}(\delta\Phi) = \nabla \log h$, and so $\delta\Phi = 0$ iff h is constant. In this case $\mathcal{X}(R^E) = \mathcal{X}(R^M)$, $p_1(R^E) = p_1(R^M)$. If M is compact and Φ is harmonic, $\Delta\Phi = 0$, then $d\Phi = 0$ and $\delta\Phi = 0$. QED

9 Homotopic almost complex structures

In this section we consider a connected compact oriented Riemannian 4-manifold (M, g) with its Levi-Civita connection and we investigate obstructions for two g -orthogonal almost complex

structures to be homotopic. This has been studied in [10], in a different context, using shadows. We will use our framework. If J_0 and J_1 are two positive g -orthogonal almost complex structures

$$J_1 = \cos \theta(p) J_0 + \tilde{H}(p) = \cos \theta(p) J_0 + \sin \theta(p) H(p) \quad (9.1)$$

where $\theta(p) \in [0, \pi]$ is the *angle* function between J_0 and J_1 , uniquely determined by the smooth function on M $\cos \theta(p) = \frac{1}{2} \langle \omega_{J_0}, \omega_{J_1} \rangle$, and $\tilde{H}(p)$ is a smooth section on E_{J_0} defined on all M . Its normalized section H may not be defined at the zero set $\mathcal{Z} = \Sigma \cup \Gamma$ of $\tilde{H}$, where $\Sigma = J_1 \cap -J_0 = \{p \in M : J_1(p) = -J_0(p)\}$ is the set of *anti-complex* points and $\Gamma = J_1 \cap J_0 = \{p \in M : J_1(p) = J_0(p)\}$ is the set of *common complex* points. Any existent homotopy from J_0 to J_1 along g -orthogonal almost complex structures must be of the form $J_t(p) = \cos \theta(p, t) J_0(p) + \tilde{H}(p, t)$. If $\Sigma = \emptyset$ then $\tilde{H}(p)$ only vanish at points p with $\theta(p) = 0$. In this case J_1 is homotopic to J_0 . For example, we may take

$$\tilde{H}(p, t) = \sqrt{\frac{t(2-t(1-\cos \theta(p)))}{(1+\cos \theta(p))}} \tilde{H}(p) =: \sin \theta(p, t) H(p) = \frac{\sin \theta(p, t)}{\sin \theta(p)} \tilde{H}(p) \quad (9.2)$$

$$\cos \theta(p, t) = 1 - t(1 - \cos \theta(p)) \quad (9.3)$$

$$J_t = \cos \theta(p, t) J_0 + \tilde{H}(p, t) = \cos \theta(p, t) J_0 + \sin \theta(p, t) H(p). \quad (9.4)$$

Note that in this case $\cos \theta(p, t)$ is smooth on $M \times [0, 1]$, $\tilde{H}$ smooth on $M \times]0, 1]$ and continuous on $M \times [0, 1]$, but not necessarily smooth, since $\frac{\sin \theta(p, t)}{\sin \theta(p)}$ may only be continuous at $t = 0$. If we reparameterize t on (9.2) and (9.3) by a smooth function $\tau(t)$ with $\tau(t) = e^{-\frac{2}{t}}$, $\tau(0) = 0$ on a neighbourhood of $t = 0$, then J_t becomes smooth at $t = 0$. Note also that $\sin \theta(p, t) = \frac{\|\tilde{H}\|}{\sqrt{2}}$ (norm in Λ_+^2) may only be continuous for $p \in \Gamma \cup \Sigma$. Hence, we have:

Proposition 9.1 *Any two positive g -orthogonal almost complex structures, everywhere linearly independent, except at common complex points, are homotopic.*

Nevertheless, for generic J_1 such set Σ is a smooth surface. In fact (see [10]) any almost complex structure J can be seen as an embedded 4-submanifold $J(M)$ into the 6-dimensional manifold $U\Lambda_+^2 TM$, the total space of the sphere bundle (of radius $\sqrt{2}$) of $\Lambda_+^2 TM$, by $p \rightarrow (p, \omega_J(p))$. Since Σ is the intersection set of $J_1(M)$ with $-J_0(M)$, this is a surface for generic J_1 . Moreover, an orientation, depending on J_1 and J_0 , can be given on each connected component of Σ ([10]).

Σ can be seen as an obstruction to the existence of a homotopy from J_0 to J_1 , but we can always consider a smooth family of g -orthogonal almost complex structures J_t , defined away from Σ as in (9.4). Then we consider an almost complex structure $\tilde{J}$ on $\pi^{-1}T(M \sim \Sigma)$, where $\pi : M \times [0, 1] \rightarrow M$, $\pi(p, t) = p$, defined by $\tilde{J}_{(p, t)} = (J_t)_p$. Let $\tilde{\omega}_{(p, t)} = (\omega_{J_t})_p = \omega_{\tilde{J}}$, and define the bundle $\tilde{E} = E_{\tilde{J}}$ by $\tilde{E}_{(p, t)} = (E_{J_t})_p =: (E_t)_p$. Then $\pi^{-1}\Lambda_+^2 TM = \mathbf{R}\tilde{\omega} \oplus \tilde{E}$. For $X \in T_p M$ we have $\nabla_{(X, 0)}^{\pi^{-1}} \omega_{\tilde{J}}(p, t) = \nabla_X \omega_{J_t}$, where $\nabla^{\pi^{-1}}$ is the pullback connection on $\pi^{-1}\Lambda_+^2 TM$, and for f a function on $M \times [0, 1]$ and s a section of $\Lambda_+^2 TM$, $\nabla_{(X, 0)}^{\tilde{E}}(f\pi^{-1}s)(p, t) = \nabla_X^{E_t}(f(\cdot, t)s)$. Moreover $R^+(\frac{d}{dt}, (X, 0)) = R^{M \times [0, 1]}(\frac{d}{dt}, (X, 0)) = 0$, where now ∇^+ denotes the induced connection on $\pi^{-1}\Lambda_+^2 TM$ and R^+ is its curvature tensor. Similar equations to (3.8) and (3.9) hold replacing J by $\tilde{J}$, and one has for $X, Y \in T_p M$, $\eta_{\tilde{J}}(p, t)(X, Y) = \eta_{J_t}(p)(X, Y)$. Therefore

$$c_1(\tilde{E})_{(p, t)}((X, 0), (Y, 0)) = c_1(E_t)_p(X, Y), \quad (9.5)$$

$$c_1(\tilde{E})_{(p, t)}(\frac{d}{dt}, (X, 0)) = \frac{1}{2\pi} \eta_{\tilde{J}}(p, t)(\frac{d}{dt}, (X, 0)). \quad (9.6)$$

So we have a closed 2-form $c_1(\tilde{E})$ on $(M \sim \Sigma) \times [0, 1]$ with $i_t^* c_1(\tilde{E}) = c_1(E_t)$ where $i_t(p) = (p, t)$. From Prop. 3.1 and remark 4 of section 7 we have proved a first conclusion:

Proposition 9.2 *If J_0 and J_1 are two positive homotopic g -orthogonal almost complex structures on M , then as real cohomology classes $c_1(M, J_0) = c_1(M, J_1)$. In particular, if $H^2(M; \mathbb{Z})$ has no torsion (for example M is simply connected) then E_{J_0} and E_{J_1} are isomorphic complex line bundles over M .*

At $p \notin \Sigma \cup \Gamma$ we define, $\omega_2(0)_p = H(p)$, $\omega_3(0)_p = J_0(p)H(p)$, $\omega_1(0)_p = \omega_0$. Then $\omega_1(t)_p := \tilde{\omega}(p, t) = \cos \theta(p, t)\omega_1(0)_p + \sin \theta(p, t)\omega_2(0)_p$, and a d.o.n.b. (of norm $\sqrt{2}$) of $\tilde{E}_{(p,t)}$ is given by

$$\omega_2(t)_p = -\sin \theta(p, t)\omega_1(0)_p + \cos \theta(p, t)\omega_2(0)_p, \quad \omega_3(t)_p = \omega_3(0)_p.$$

Thus,

$$\begin{aligned} \nabla_{(X,0)}^+ \tilde{\omega}(p, t) &= \cos \theta(p, t) \nabla_X^+ \omega_1(0)_p + \sin \theta(p, t) \nabla_X^+ \omega_2(0)_p \\ &\quad + d \cos \theta(p, t) (X, 0) \omega_1(0)_p + d \sin \theta(p, t) (X, 0) \omega_2(0)_p \\ \nabla_{\frac{d}{dt}}^+ \tilde{\omega}(p, t) &= \frac{d}{dt} \cos \theta(p, t) \omega_1(0)_p + \frac{d}{dt} \sin \theta(p, t) \omega_2(0)_p \end{aligned}$$

$$\begin{aligned} \eta_{\tilde{J}}\left(\frac{d}{dt}, (X, 0)\right) &= \frac{1}{2} \text{Vol}_{\tilde{E}} \left(\nabla_{\frac{d}{dt}}^+ \tilde{\omega}, \nabla_{(X,0)}^+ \tilde{\omega}(p, t) \right) = \\ &= \left(-\sin \theta(p, t) \frac{d}{dt} \cos \theta(p, t) + \cos \theta(p, t) \frac{d}{dt} \sin \theta(p, t) \right) \\ &\quad \cdot \left(\cos \theta(p, t) \langle \nabla_X J_0, J_0 H(p) \rangle + \sin \theta(p, t) \langle \nabla_X H, J_0 H(p) \rangle \right) \end{aligned}$$

where $\nabla_X J$ is the covariant derivative in $\text{Skew}(TM)$ and $\langle \cdot, \cdot \rangle$ is half the usual Riemannian metric on that vector bundle (that corresponds to the Riemannian metric of $\Lambda^2 TM$). For simplicity from now on we will denote by ∇ the connection ∇^+ of $\Lambda_+^2 TM$. Note that

$$\begin{aligned} -\cos \theta \sin \theta \frac{d}{dt} \cos \theta + \cos^2 \theta \frac{d}{dt} \sin \theta &= -\frac{1}{2} \sin \theta \frac{d}{dt} \cos^2 \theta + \cos^2 \theta \frac{d}{dt} \sin \theta \\ &= \frac{1}{2} \sin \theta \frac{d}{dt} \sin^2 \theta + \cos^2 \theta \frac{d}{dt} \sin \theta = \sin^2 \theta \frac{d}{dt} \sin \theta + \cos^2 \theta \frac{d}{dt} \sin \theta = \frac{d}{dt} \sin \theta. \end{aligned}$$

Similarly $\cos \theta \sin \theta \frac{d}{dt} \sin \theta - \sin^2 \theta \frac{d}{dt} \cos \theta = -\frac{d}{dt} \cos \theta$. Consequently

$$c_1(\tilde{E})_{(p,t)}\left(\frac{d}{dt}, (X, 0)\right) = \frac{1}{2\pi} \left(-\frac{d}{dt} \cos \theta \langle \nabla_X H, J_0 H \rangle + \frac{d}{dt} \sin \theta \langle \nabla_X J_0, J_0 H \rangle \right) \quad (9.7)$$

and we have got the following formula:

Proposition 9.3 *If J_0 and J_1 are two g -orthogonal almost complex structures on M , with $J_1 = \cos \theta J_0 + \tilde{H}$ where $\tilde{H} \in C^\infty(E_{J_0})$, then*

$$4\pi(c_1(M, J_1) - c_1(M, J_0)) = d\tilde{T} + dG \quad (9.8)$$

where $\tilde{T}$ and G are 1-forms on M , G globally defined on M and $\tilde{T}$ defined away from Σ by

$$\forall X \in T_p M \quad \tilde{T}(X) = \frac{1}{(1 + \cos \theta(p))} \langle \nabla_X \tilde{H}, J_0 \tilde{H} \rangle, \quad G(X) = \langle \nabla_X J_0, J_0 \tilde{H} \rangle. \quad (9.9)$$

where now $\langle \cdot, \cdot \rangle$ is the usual inner product of $\text{Skew}(TM) \subset TM^* \otimes TM$ (twice the one of $\Lambda_+^2 TM$).

Proof. We apply remark 4 of section 7 to $c_1(\tilde{E})$ given by (9.5), (9.6), and satisfying (9.7), smooth on $(M \sim \Sigma \cup \Gamma) \times [0, 1]$, obtaining $T(X)$ defined for $p \notin \Sigma \cup \Gamma$,

$$\begin{aligned} T(X) &= 4\pi \int_0^1 c_1(\tilde{E})\left(\frac{d}{dt}, (X, 0)\right) dt = \int_0^1 -\frac{d}{dt} \cos \theta(p, t) \langle \nabla_X H, J_0 H \rangle + \frac{d}{dt} \sin \theta(p, t) \langle \nabla_X J_0, J_0 H \rangle \\ &= (1 - \cos \theta(p)) \langle \nabla_X H, J_0 H \rangle + \sin \theta(p) \langle \nabla_X J_0, J_0 H \rangle = \tilde{T}(X) + G(X). \end{aligned}$$

T can be smoothly extended to Γ . $\tilde{T}$ may not be defined at Σ , but G is smooth on all M .
QED

Corollary 9.1

$$\int_M c_1(M, J_1) \wedge c_1(M, J_0) = \int_M p_1(\Lambda_+^2 TM) - \frac{1}{32\pi^2} \int_M d(\tilde{T} \wedge d(\tilde{T} + 2G)).$$

Proof. From Prop.3.1. $16\pi^2(c_1(M, J_1) - c_1(M, J_0))^2 = 16\pi^2(2p_1(\Lambda_+^2 TM) - 2c_1(M, J_1) \wedge c_1(M, J_0))$. Now formula of Prop. 9.3 gives

$$c_1(M, J_1) \wedge c_1(M, J_0) = p_1(\Lambda_+^2 TM) - \frac{1}{32\pi^2} d(\tilde{T} \wedge d\tilde{T} + G \wedge dG + 2\tilde{T} \wedge dG). \quad \text{QED}$$

Proof of Theorem 1.2. Recall that ω_0 is a closed 2-form. From Prop.9.3

$$4\pi(c_1(M, J_1) - c_1(M, J_0)) \wedge \omega_0 = d\tilde{T} \wedge \omega_0 + dG \wedge \omega_0 = d(\tilde{T} \wedge \omega_0) + d(G \wedge \omega_0).$$

Since $\tilde{T} \wedge \omega_0$ is a 3-form, $d(\tilde{T} \wedge \omega_0) = -\text{div}((\star(\tilde{T} \wedge \omega_0))^\sharp) \text{Vol}_M$, where $\star : \Lambda^3 TM^* \rightarrow TM^*$ is the star operator. Now $\omega_0 = e_*^{12} + e_*^{34}$ where $e_2 = J_0 e_1$, $e_4 = J_0 e_3$, and so

$$\begin{aligned} \star(\tilde{T} \wedge \omega_0) &= \star(\tilde{T}(e_3)e_*^{312} + \tilde{T}(e_4)e_*^{412} + \tilde{T}(e_1)e_*^{134} + \tilde{T}(e_2)e_*^{234}) \\ &= -\tilde{T}(J_0 e_4)e_*^4 - \tilde{T}(J_0 e_3)e_*^3 - \tilde{T}(J_0 e_2)e_*^2 - \tilde{T}(J_0 e_1)e_*^1 = -\tilde{T}J_0. \end{aligned}$$

Consequently, $4\pi(c_1(M, J_1) - c_1(M, J_0)) \wedge \omega_0 = \text{div}((\tilde{T}J_0)^\sharp) \text{Vol}_M + d(G \wedge \omega_0)$. Integration over M and Stokes on the second term leads to the first equality of Theorem 1.2. Assume $\nabla^{E_{J_0}} \tilde{H}$ is anti- J_0 -complex, that is $\nabla_{J_0 X}^{E_{J_0}} \tilde{H} = -J_0 \nabla_X^{E_{J_0}} \tilde{H}$. Then,

$$(1 + \cos \theta) \tilde{T}(J_0(X)) = \langle \nabla_{J_0 X} \tilde{H}, J_0 \tilde{H} \rangle = \langle \nabla_{J_0 X}^{E_{J_0}} \tilde{H}, J_0 \tilde{H} \rangle = -\langle \nabla_X \tilde{H}, \tilde{H} \rangle.$$

We have $\tilde{H} = \sin \theta H$ and $\langle \nabla_X H, H \rangle = 0$. Thus, $\langle \nabla_X \tilde{H}, \tilde{H} \rangle = \frac{1}{2} d \sin^2 \theta(X)$, and so, $\text{div}((\tilde{T}J_0)^\sharp) = -\frac{1}{2} \text{div}\left(\frac{\nabla \sin^2 \theta}{(1 + \cos \theta)}\right)$. Now $\frac{\nabla \sin^2 \theta}{(1 + \cos \theta)} = \frac{2 \sin \theta \cos \theta \nabla \theta}{1 + \cos \theta}$, and

$$\begin{aligned} \frac{2 \sin \theta \cos \theta}{1 + \cos \theta} &= \frac{4 \sin \frac{\theta}{2} \cos \frac{\theta}{2} \cos \theta}{2 \cos^2 \frac{\theta}{2}} = 2 \tan \frac{\theta}{2} \cos \theta = (2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}) \frac{\cos \theta}{\cos^2 \frac{\theta}{2}} = (2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}) \left(2 - \frac{1}{\cos^2 \frac{\theta}{2}}\right) \\ &= \left(4 \sin \frac{\theta}{2} \cos \frac{\theta}{2} - 2 \frac{\sin \frac{\theta}{2}}{\cos \frac{\theta}{2}}\right) = \left(2 \sin \theta - 2 \tan \frac{\theta}{2}\right) = \left(2 \sin \theta - \frac{2 \sin \frac{\theta}{2}}{\cos \frac{\theta}{2}}\right) \end{aligned}$$

That is $\frac{\nabla \sin^2 \theta}{(1 + \cos \theta)} = \nabla(-2 \cos \theta + 4 \log |\cos \frac{\theta}{2}|)$, where $\theta \in [0, \pi]$, obtaining

$$\text{div}((\tilde{T}J_0)^\sharp) = -\frac{1}{2} \text{div}\left(\frac{\nabla \sin^2 \theta}{(1 + \cos \theta)}\right) = \Delta\left(\cos \theta - 2 \log(\cos \frac{\theta}{2})\right).$$

Since $\log(\cos \frac{\theta}{2}) = \frac{1}{2} \log\left(\frac{1+\cos \theta}{2}\right) = \frac{1}{2}(\log(1+\cos \theta) - \log(2))$, integration over M of $\text{div}((\tilde{T}J_0)^\#)$ and Stokes on the first term $\Delta \cos \theta$ gives formula (1.4) in the Theorem.

Now assume Σ is a finite disjoint union of connected compact and orientable submanifolds Σ_i of dimension $d_i \leq 2$. Then for a sufficiently small tubular neighbourhood of Σ , $V(\Sigma) = \bigcup_i V(\Sigma_i)$, $\partial(M \sim V(\Sigma)) = \bigcup_i -\partial V(\Sigma_i)$. Below we specify such tubular neighbourhood. For notational simplicity we assume $\Sigma = \Sigma_i$.

For each $r > 0$, let G_r and C_r be the subsets of the total space $N\Sigma$ of the normal bundle $T\Sigma^\perp$ of Σ in M given by

$$G_r = \{(p, w) : p \in \Sigma, w \in T_p\Sigma^\perp, \|w\| \leq r\}, \quad C_r = \{(p, w) : p \in \Sigma, w \in T_p\Sigma^\perp, \|w\| = r\}$$

and for r sufficiently small (say $r \leq r_0 \leq 1$) the exponential map of M restricted to G_r , denoted by $\exp : G_r \rightarrow M$, $\exp(p, w) = \exp_p(w)$ defines a diffeomorphism onto $V(\Sigma, r) = \{q \in M : d(q, \Sigma) \leq r\}$ and $C(\Sigma, r) = \{q \in M : d(q, \Sigma) = r\}$ is its boundary. Let $S(p, r)$ denote the sphere of radius r in $T_p\Sigma^\perp$, $\sigma(q) = d(q, \Sigma)$, and for each $w \in T_p\Sigma^\perp$, $\gamma_{(p,w)}(r) = \exp_p(rw)$ is the geodesic normal to Σ , starting at p with initial velocity w . Thus, $s(p, w) := \sigma(\exp(p, w)) = \|w\|$, is just the Euclidean norm in $T_p\Sigma^\perp$. Since $N\Sigma$ is the total space of a Riemannian vector bundle, then it has a natural Riemannian structure such that $\pi : N\Sigma \rightarrow \Sigma$ is a Riemannian submersion. The volume element $\text{Vol}_{N\Sigma}$ for such metric satisfies $\text{Vol}_{N\Sigma}(p, w) = \text{Vol}_\Sigma(p) \wedge ds(p, w)$ and $\text{Vol}_{C_r}(p, w) = \text{Vol}_\Sigma(p) \wedge \text{Vol}_{S(p,r)}(w)$, where $r = \|w\|$. For each $u \in S(p, 1)$ define $\vartheta_u(r) = \langle \text{Vol}_{N\Sigma}(p, ru), \exp^* \text{Vol}_M(p, ru) \rangle$. This function $\vartheta_u(r)$ measures the volume distortion by $\exp$ in the direction u . It satisfies $\vartheta_u(0) = 1$ and a Riccati differential equation. We recall the following (see [14]): (1) $\nu(q) = \nabla \sigma(q)$ is the unit outward of $C(\Sigma, r)$, (2) $\nu(\gamma_{(p,u)}(r)) = \gamma'_{(p,u)}(r)$, (3) $ds \wedge *ds$ and $d\sigma \wedge *d\sigma$ are the volume elements of $N\Sigma$ and M respectively. (4) $*ds$ and $*d\sigma$ are the volume elements of each hypersurface C_r of $N\Sigma$ and $C(\Sigma, r)$ of M respectively, and $\exp^*(*d\sigma)(p, w) = \vartheta_{\frac{w}{r}}(r)(*ds)(p, w)$, where $r = \|w\|$. We have

$$\begin{aligned} \int_M -\frac{1}{4\pi} \Delta \log(1 + \cos \theta) \text{Vol}_M &= \lim_{\epsilon \rightarrow 0} \int_{M \sim V(\Sigma, \epsilon)} -\frac{1}{4\pi} \Delta \log(1 + \cos \theta) \text{Vol}_M \\ &= \lim_{\epsilon \rightarrow 0} \int_{C(\Sigma, \epsilon)} \frac{1}{4\pi} \langle \nabla(\log(1 + \cos \theta)), \nu \rangle * d\sigma = \lim_{\epsilon \rightarrow 0} \int_{C_\epsilon} \frac{1}{4\pi} (\exp)^* (d(\log(1 + \cos \theta + 1))(\nu) * d\sigma) \\ &= \lim_{\epsilon \rightarrow 0} \int_\Sigma \frac{1}{4\pi} \left(\int_{S(p, \epsilon)} d(\log(1 + \cos \theta))_{(\exp_p(w))}(\nu) \vartheta_{\frac{w}{\epsilon}}(\epsilon) d_{S(p, \epsilon)}(w) \right) d_\Sigma(p) \\ &= \lim_{\epsilon \rightarrow 0} \int_\Sigma \frac{1}{4\pi} \left(\int_{S(p, 1)} \epsilon^{3-d_i} \vartheta_u(\epsilon) \frac{d}{dr} \Big|_{r=\epsilon} \log(\phi_{(p,u)}(r)) d_{S(p, 1)}(u) \right) d_\Sigma(p) \end{aligned} \quad (9.10)$$

where $\phi_{(p,u)}(r) = (1 + \cos \theta)(\gamma_{(p,u)}(r))$. The function $\kappa(p, u)$ = the order of the zero $r = 0$ of $\phi_{(p,u)}(r)$, is defined in the total space of $N^1\Sigma$ with values in $[1, +\infty]$ and satisfies $\kappa(p', u') \leq \kappa(p, u)$ for all (p', u') in a neighbourhood of (p, u) . In particular it is upper semi-continuous, and so it is measurable. Now we have for $k = \kappa(p, u) < +\infty$ and $\forall 0 < \epsilon < r_0$

$$0 \neq \phi_{(p,u)}(\epsilon) = \epsilon^k (A(p, u) + \epsilon B(p, u, \epsilon)) \quad (9.11)$$

$$\text{where } A(p, u) = \frac{1}{k!} \frac{d^k}{dr^k} \phi_{(p,u)}(0) \neq 0, \quad B(p, u, \epsilon) = \int_0^1 \frac{(1-s)^k}{k!} \frac{d^{k+1}}{dr^{k+1}} \phi_{(p,u)}(s\epsilon) ds \quad (9.12)$$

Thus,

$$\epsilon \frac{d}{d\epsilon} \log(\phi_{(p,u)}(\epsilon)) = \epsilon \frac{\frac{d}{d\epsilon} \phi_{(p,u)}(\epsilon)}{\phi_{(p,u)}(\epsilon)} = \kappa(p, u) + \epsilon \frac{B(p, u, \epsilon) + \epsilon \frac{d}{d\epsilon} B(p, u, \epsilon)}{A(p, u) + \epsilon B(p, u, \epsilon)}. \quad (9.13)$$

Assume that for almost all $(u, p) \in N^1\Sigma$, $\kappa(p, u) < +\infty$. Then

$$(9.10) = \lim_{\epsilon \rightarrow 0} \frac{1}{4\pi} \int_{\Sigma} \int_{S(p,1)} \epsilon^{2-d_i} \vartheta_u(\epsilon) \left(\kappa(p, u) + \frac{\epsilon(B(p, u, \epsilon) + \epsilon \frac{d}{d\epsilon} B(p, u, \epsilon))}{A(p, u) + \epsilon B(p, u, \epsilon)} \right) d_{S(p,1)} d_{\Sigma}(p) \quad (9.14)$$

and for almost all (u, p)

$$\lim_{\epsilon \rightarrow 0} \vartheta_u(\epsilon) \left(\kappa(p, u) + \frac{\epsilon(B(p, u, \epsilon) + \epsilon \frac{d}{d\epsilon} B(p, u, \epsilon))}{A(p, u) + \epsilon B(p, u, \epsilon)} \right) = \vartheta_u(0) \kappa(p, u) = \kappa(p, u).$$

In order to interchange $\lim_{\epsilon \rightarrow 0}$ with $\int_{\Sigma} \int_{S(p,1)}$ we need some conditions to be satisfied. Under the assumption of controlled zero set, and if $d_i \leq 2$, we may apply the dominate convergence theorem to (9.14), and only the terms with $d_i = 2$ do not vanish. The last inequality follows immediately from (3.7). QED

Proposition 9.4 *Assume $\Sigma = \cup \Sigma_i$ as in Theorem 1.2. Then $(1 + \cos \theta)$ has controlled zero set if (1), or (2) holds:*

- (1) $\kappa \in L^1(N^1\Sigma)$ and there exist a constant $r_0 > 0$, and non-negative functions h, d on $N^1\Sigma$ with $h/d \in L^1(N^1\Sigma)$ such that for almost all $(p, u) \in N^1\Sigma$, $|A(p, u)| \geq \sup_{r \leq r_0} |\frac{r}{k!} \frac{d^{k+1}}{dr^{k+1}} \phi_{(p,u)}(r)| + d(p, u)$ and $\sup_{r \leq r_0} \left\{ |\frac{r}{k!} \frac{d^{k+1}}{dr^{k+1}} \phi_{(p,u)}(r)|, |\frac{r^2}{k!} \frac{d^{k+2}}{dr^{k+2}} \phi_{(p,u)}(r)| \right\} \leq h(p, u)$, where $k = \kappa(p, u)$.
- (2) For all (p, u) , $\phi_{(p,u)}(r)$ is polynomial on $r \in [0, r_0]$ with coefficient $A(p, u)$ uniformly bounded away from 0, i.e. $|A(p, u)| \geq c \forall (p, u)$, where $c > 0$ is constant.

Proof. (1) For a.e. (p, u) , and for $0 < \epsilon < r_0$, $|A(p, u) + \epsilon B(p, u, \epsilon)| \geq |A(p, u)| - \epsilon |B(p, u, \epsilon)| \geq |A(p, u)| - \sup_{r \leq r_0} |\frac{r}{k!} \frac{d^{k+1}}{dr^{k+1}} \phi_{(p,u)}(r)| \geq d(p, u)$. Moreover $\epsilon |B(p, u, \epsilon) + \epsilon \frac{d}{d\epsilon} B(p, u, \epsilon)| \leq h(p, u)$. Therefore $|(9.13)| \leq \kappa(p, u) + \frac{h(p, u)}{d(p, u)}$, what implies that $(1 + \cos \theta)$ has a controlled zero set. (2) Let $r_0 > 0$ sufficiently small s.t. all geodesics $\gamma_{(p,u)}(\epsilon)$ are defined for $0 \leq \epsilon < r_0$. In this case κ takes only finite values, and since it is u.s.c., it has a maximum k_0 in $N\Sigma^1$. Then we may take $C := \sum_{\mu \leq k_0} \max_{(p,u) \in N\Sigma^1, \epsilon \leq \epsilon_0} |\frac{1}{(\mu+1)!} \frac{d^{\mu+1}}{dr^{\mu+1}} \phi_{(p,u)}(\epsilon)|$. Then we can find $\epsilon_0 > 0$ s.t. $|A(p, u) + \epsilon B(p, u, \epsilon)| \geq \frac{c}{2}$ for all $\epsilon \leq \epsilon_0$ and all (p, u) . QED

Corollary 9.2 *If J_0 and J_1 satisfy the conditions of the last part of Theorem 1.2, then:*

- (1) $\left(\int_M \|c_1(M, J_1)\|^2 Vol_M \right)^{\frac{1}{2}} \geq \frac{1}{4\sqrt{2}\pi(Vol(M))^{\frac{1}{2}}} \int_M (s^M + \frac{1}{2} \|\nabla \omega_0\|^2) Vol_M$, with equality iff $c_1(M, J_1) = f\omega_0$ for some smooth function $f : M \rightarrow \mathbf{R}$.
- (2) If, $c_1(M, J_1)$ is L^2 -orthogonal to ω_0 then $\int_M s^M Vol_M \leq 0$ and $\left| \int_M s_{J_0}^M Vol_M \right| \leq - \int_M s^M Vol_M$. If $\int_M s^M Vol_M = 0$ then J_0 is Kähler.

Proof. From last inequalities of Theorem 1.2 we clearly obtain (1). Moreover, we have $0 \geq \int_M (s^M + \frac{1}{2} \|\nabla \omega_0\|^2) Vol_M$. In particular $\int_M s^M Vol_M \leq 0$, and using the equality (3.6), and that $s^M \leq s_{J_0}^M$, we obtain the proof of (2). QED

Remark 5. If M is a surface and $f : M \rightarrow \mathbf{R}$ is a function of absolute value type at a zero p

([11]), i.e. locally $|f(z)| = |z|^k \psi(z)$ where z is an isothermal coordinate around p , and $\psi(z) > 0$, then $\kappa(p, u) = k$ for all u . We also say that p is a zero of homogeneous order k .

Remark 6. Assume J_0 is almost Kähler. Note that if J_0 is Kähler, E_{J_0} is a parallel subbundle of $\bigwedge_+^2 TM$. If $\tilde{H}$ is a non-zero parallel section of $\bigwedge_+^2 TM$ then by lemma 3.1 M is hyper-Kähler, with respect to $(J_0, J_1 = H)$. If $\tilde{H}$ is a parallel section of E_{J_0} , then J_1 is Kähler iff J_0 is so.

Proof of Proposition 1.2. Since J_0 is Kähler, ω_0 is parallel and so harmonic, and the Weitzenböck operator satisfies $A(\omega_0) = 0$ (see (3.2)). Since J_1 is almost Kähler, then ω_1 is harmonic. Polarization of the Weitzenböck formula (second equality of (3.3)) and using the fact that A is a symmetric operator, gives

$$\Delta^+(\langle \omega_0, \omega_1 \rangle) = -\langle \Delta \omega_0, \omega_1 \rangle - \langle \omega_0, \Delta \omega_1 \rangle + 2\langle \nabla \omega_0, \nabla \omega_1 \rangle + \langle A(\omega_0), \omega_1 \rangle + \langle \omega_0, A(\omega_1) \rangle = 0$$

that is, $\cos \theta$ is an harmonic function. Since M is closed $\cos \theta$ is constant. If $\cos \theta \neq \pm 1$ Prop. 9.1 gives the homotopy, and orthonormalization of Gram-Schmidt of ω_0, ω_1 gives two self-dual 2-forms ω_0 and ω'_1 that correspond to two g -orthogonal complex structures J_0 , Kähler, and J_1 , almost Kähler, that anti-commute. Lemma 3.1 proves Proposition 1.2. QED

10 Applications

10.1 Surfaces

(1) Let (M, g) be a compact Riemann surface of Gauss curvature K . If $\xi : U \subset \mathbf{C} \rightarrow M$ is a conformal coordinate with $g = \lambda^2 \langle, \rangle$ where λ^2 is the conformal factor, then $K = -\frac{1}{2} \Delta \log \lambda^2$ and the Euler number of M is $\mathcal{X}(M) = \frac{1}{2\pi} \int_M K Vol_g$. If we take a conformal change of the metric $\hat{g} = hg$ where h is a non-negative smooth real function, then $Vol_{\hat{g}} = h Vol_g$, and ξ is still a conformal coordinate for $\hat{M} = (M, \hat{g})$ with conformal factor $h\lambda^2$. Hence

$$\begin{aligned} \hat{K} &= -\frac{1}{2} \hat{\Delta} \log h \lambda^2 = -\frac{1}{2} (h^{-1} \Delta \log h + h^{-1} \Delta \log \lambda^2) = -\frac{1}{2} h^{-1} \Delta \log h + h^{-1} K \\ \mathcal{X}(\hat{M}) &= \mathcal{X}(M) - \frac{1}{2\pi} \int_M \frac{1}{2} \Delta \log h Vol_g \end{aligned} \quad (10.1)$$

If on a isothermal coordinate of each zero or infinity of h , h is of the form $h(z) = |z|^k \psi(z)$ where $\psi \neq 0, \infty$ is a smooth function and $k \in \mathbf{Z}$, h is said to be of *absolute value type* at those points (see remark 5). In this case

$$-\frac{1}{2\pi} \int_M \Delta \log h Vol_g = \text{number of zeros of } h - \text{number of infinities of } h \quad (10.2)$$

where the zeros or infinities of h are counting including multiplicity (see [11]).

(2) Consider $\sigma : S^2 \rightarrow \mathbf{R}^2$ the stereographic projection, $\phi(x_0, x_1, x_2) = \omega = \frac{1}{1-x_0}(x_1, x_2)$, then σ defines an isometry between $S^2 \sim (1, 0, 0)$ with the metric hg_S and $\mathbf{R}^2$ with the Euclidean metric g_0 , where $h = \frac{1}{(1-x_0)^2}$ and g_S is the usual metric of S^2 . Since $(\mathbf{R}^2, g_0)$ is a flat space, its Euler form is a zero form, and so $0 = \mathcal{X}(\mathbf{R}^2) = \mathcal{X}(S^2) - \frac{1}{2}(\text{number of infinities of } h)$. Note that $\frac{1}{h}(x_0, x_1, x_2) = (1-x_0)^2$ has a zero of order 4 at $(1, 0, 0)$ as a function on S^2 . We could also consider σ as an isometry between (S^2, g_S) and $\mathbf{R}^2$ with the metric $\hat{h}g_0$ where $\hat{h} = \frac{4}{(1+\|w\|^2)^2}$

and observe that ∞ is a zero of 4-order of $\tilde{h}$. Therefore $\mathcal{X}(S^2) = 2$.

(3) Let $F : M^2 \rightarrow N^4$ be a minimal immersed surface into a Kähler complex surface (N, J, g) and denote by θ the Kähler angle of F , that is $F^*\omega = \cos \theta \text{Vol}_M$ where $\omega(X, Y) = g(JX, Y)$. Let $\hat{g} = \sin^2 \theta g_M$ where g_M is the induced metric on M by F . The bundle map $\Phi : TM \rightarrow NM$, $\Phi(X) = (JX)^\perp$, where NM is the normal bundle, is a conformal bundle map with $\|\Phi(X)\|^2 = \sin^2 \theta \|X\|^2$. If T' is the torsion of the $\hat{g}$ -Riemannian connection ∇' of M that makes Φ a parallel isometry from $(TM, \hat{g}, \nabla')$ to NM with its usual metric and connection $\nabla^\perp$, then (see [24](3.2))

$$\Phi(T'(X, Y)) = \cos \theta (\nabla dF(J_\omega X, Y) - \nabla dF(X, J_\omega Y))$$

where J_ω is a g_M -orthogonal complex structure on M . Minimality of F is equivalent to the vanishing of the r.h.s. of the above equation. This means that $T' = 0$, and so ∇' is the Levi-Civita connection for $\hat{M} = (M, \hat{g})$. Moreover, one can prove that if F is not a complex submanifold, minimality of F implies $\sin^2 \theta$ is of absolute value type at its zeros and so zeros are isolated and of finite order. Taking into consideration that Φ is anti-holomorphic for the g -orthogonal complex structures on TM and NM defined by the usual orientation (and so it is a reverse orientation bundle map), (10.1) and (10.2) gives

$$-\mathcal{X}(NM) = \mathcal{X}(R') = \mathcal{X}(\hat{M}) = \mathcal{X}(M) + \sum_{p \in \mathcal{C}^+} \text{order}(p) + \sum_{p \in \mathcal{C}^-} \text{order}(p)$$

where $\mathcal{C}^+$ and $\mathcal{C}^-$ are respectively the sets of complex and anti-complex points, zeros of $\sin^2 \theta = (1 - \cos \theta)(1 + \cos \theta)$. This formula was obtained by [11], [26] and [27], that we give here a short proof, using a singular connection.

10.2 Minimal 4-manifolds

Let $F : M \rightarrow N$ be a 4-dimensional immersed Cayley submanifold M of a Calabi-Yau manifold N of complex dimension 4. This means that M is calibrated by one of the Cayley calibrations of N (see e.g. [19]), and this is equivalent to M to be minimal and with equal Kähler angles. In this case the morphism $\Phi : TM \rightarrow NM$, $\Phi(X) = (JX)^\perp$, where NM is the normal bundle, has the property of being a conformal one, with coefficient of conformality $\sin^2 \theta$, where θ is the common Kähler angle of M . For such submanifolds, the complex points are zeros of finite order of $\sin^2 \theta$. This is treated in [23] where using Theorem 1.1 of the present paper one describes the set of complex points of M as a residue formula involving the Euler (or equivalently, the Pontrjagin) numbers of M and of the normal bundle. The morphism Φ defines a singular metric $\hat{g} = \sin^2 \theta g_M$ on M , vanishing at complex points, and a connection ∇' that is $\hat{g}$ -Riemannian and has torsion and identifies $p_1(NM)$ with $p_1(R')$, and $\mathcal{X}(NM)$ with $\mathcal{X}(R')$ as in the surface case.

10.3 Isolated pole in dimension 4

Let (M, g) be a closed Riemannian 4-manifold, $p_0 \in M$, and h a smooth function on $M \sim \{p_0\}$ with $+\infty > h(p) > 0$ and $h(p_0) = 0$ or $h(p_0) = +\infty$. Assume first that g is flat on a neighbourhood of p_0 , and on a geodesic ball $B(p_0, \epsilon) \equiv B(0, \epsilon)$, $h(x) = \|x\|^{2k}$ for some integer k . Let ν be the outward unit of the (geodesic) sphere $S(p_0, \epsilon)$ of M of radius ϵ . Let $f = \log h =$

$2k \log \|x\|$. Then $\int_M \text{div}_g(P(\nabla f)) \text{Vol}_M = \lim_{\epsilon \rightarrow 0} - \int_{S(p_0, \epsilon)} g(P(\nabla f)(x), \nu(x)) \text{Vol}_{S(p_0, \epsilon)}$, where $P(\nabla f)$ is defined in (1.1). For each $x \in S(0, \epsilon)$

$$g(P(\nabla f)(x), \nu(x)) = (2k)^2(2k+6)\epsilon^{-3} \quad (10.3)$$

and so for any ϵ we obtain the integer

$$- \frac{1}{32\pi^2} \int_{S(p_0, \epsilon)} g(P(\nabla f), \nu) \text{Vol}_{S(0, \epsilon)} = -\frac{1}{2}k^2(k+3). \quad (10.4)$$

In the general case, if (M, g) is not flat near p_0 but on a geodesic ball of p_0 , $f = \log(\tilde{f}h)$ where $+\infty > \tilde{f} > 0$ is smooth and $h(\exp_{p_0}(w)) = \|w\|^{2k}$ (e^f is of "absolute value type" at p_0), using normal coordinates at p_0 we can easily compute $g(P(\nabla f), \nu)$ on a geodesic ball of radius ϵ , and taking the limit $\epsilon \rightarrow 0$ of the l.h.s. of (10.4) we obtain the same integer on r.h.s. of (10.4).

11 Quaternionic-Hermitian 8-manifolds

If (M^8, Q, g) is an oriented almost quaternionic Hermitian (AQH) 8-manifold (see [2] for definitions), its fundamental form Ω is a self-dual 4-form of norm $\sqrt{\frac{10}{3}}$. If M is quaternionic Kähler (QK) then the cohomology class $[\Omega]$ of the fundamental 4-form represents the first Pontrjagin class of E (see [20], [8]), where $E = \text{span}\{J_1, J_2, J_3 = J_1 J_2\}$ is the rank-3 bundle locally defined by a pair of anti-commuting g -orthogonal positive almost complex structures J_1, J_2 of Q . If M is only almost QK (AQK) we may compare $[\Omega]$ with $p_1(E)$, and we may also use the Obata connection to describe characteristic classes, in a similar way for almost complex structures. For each pair Q_0 and Q_1 of AQH structures on M , with fundamental forms Ω_0 and Ω_1 , we define at each point $p \in M$ an angle $\theta(p) \in [0, \pi]$ as $\cos \theta(p) = \frac{3}{10} \langle \Omega_0(p), \Omega_1(p) \rangle$, where $\langle, \rangle$ is the Hilbert-Schmidt inner product. Then it is defined a smooth section $\tilde{H}$ of $\Lambda_+^4(TM)$, orthogonal to Ω_0 and with $\|\tilde{H}\|^2 = \frac{3}{10} \sin^2 \theta$, given by

$$\Omega_1 = \cos \theta(p) \Omega_0 + \tilde{H}(p). \quad (11.1)$$

If Q_0 and Q_1 are both AQK we would like to compare the cohomology classes of Ω_i or compare $p_1(E_0)$ with $p_1(E_1)$, in terms of a PDE on θ , and find conditions to conclude: (a) θ is constant; or (b) Q_1 is QK; (c) Ω_1 and Ω_0 are homotopic; or (d) E_1 and E_2 are isomorphic. As in section 9, taking $\cos \theta(p, t)$ and $H(\tilde{p}, t)$ defined for instance as in (9.3) and (9.2), and conveniently reparameterizing t near $t = 0$, we have a simple answer to (c):

Proposition 11.1 *If Q_0 and Q_1 are two AQH structures with no anti-quaternionic points, i.e. points with $\cos \theta(p) = -1$, then a homotopy Ω_t of self-dual 4-forms exist from Ω_0 to Ω_1 . Furthermore, if Q_0 and Q_1 are AQK and $p \rightarrow \cos \theta(p)$ is constant, then the homotopy can be taken by closed self-dual 4-forms.*

If $\cos \theta(p)$ is constant, then it is also $\cos \theta(p, t)$ defined by (9.3). Clearly it follows that $\Omega_t = \cos \theta(p, t) \Omega_0 + \tilde{H}(p, t)$ is also closed. So, existence of anti-quaternionic points can be an obstruction to homotopy, but the non-existence of them does not guarantee that we can take Ω_t a fundamental form for some AQH structure, not even locally.

Weitzenböck formula (3.2) and second equality of (3.3) also hold for 4-forms, where $A = A(R^M) : \wedge^4 TM^* \rightarrow \wedge^4 TM^*$ is also symmetric for the Hilbert-Schmidt inner product, as we see next. Denote by $\overline{R^M}$ the induced curvature tensor of $\wedge^4 TM^*$: if Ω is a 4-form, $\overline{R^M}(X, Y)\Omega(X_1, \dots, X_4) = \sum_{1 \leq j \leq 4} -\Omega(X_1, \dots, R^M(X, Y)X_j, \dots, X_4)$. Let (e_α) be an o.n. basis of $T_p M$. The Weitzenböck operator applied to Ω is given by

$$A(\Omega)(X_1, X_2, X_3, X_4) = \sum_{1 \leq \alpha \leq 8} \sum_{1 \leq k \leq 4} (-1)^k (\overline{R^M}(e_\alpha, X_k)\Omega)(e_\alpha, X_1, \dots, X_{\hat{k}}, \dots, X_4).$$

Lemma 11.1 *For any distinct $\alpha, \beta, \gamma, \mu, \nu, \sigma, \rho, \eta$:*

$$\begin{aligned} \langle A(e_*^{\alpha\beta\gamma\mu}), e_*^{\nu\sigma\rho\eta} \rangle &= \langle A(e_*^{\alpha\beta\gamma\mu}), e_*^{\alpha\sigma\rho\eta} \rangle = 0 \\ \langle A(e_*^{\alpha\beta\gamma\mu}), e_*^{\alpha\beta\rho\eta} \rangle &= 2R^M(e_\gamma, e_\rho, e_\eta, e_\mu) - 2R^M(e_\mu, e_\rho, e_\eta, e_\gamma) = -2R^M(e_\gamma, e_\mu, e_\rho, e_\eta) \\ \langle A(e_*^{\alpha\beta\gamma\mu}), e_*^{\alpha\beta\gamma\eta} \rangle &= -2R^M(e_\alpha, e_\mu, e_\alpha, e_\eta) - 2R^M(e_\beta, e_\mu, e_\beta, e_\eta) - 2R^M(e_\gamma, e_\mu, e_\gamma, e_\eta) - \text{Ricci}^M(e_\mu, e_\eta) \end{aligned}$$

and so $\langle A(\Omega), \Omega' \rangle = \langle \Omega, A(\Omega') \rangle$ for any Ω, Ω' .

Note that if Q_1 is AKQ, since Ω_1 is self-dual and closed then it is co-closed and so it is an harmonic 4-form. With similar proof to the one of Proposition 1.2 (section 9), and using Lemma 11.1, we have the following conclusion:

Proposition 11.2 *If Ω_0 is QK, Ω_1 is AQK and M is compact, then θ is constant. If $\cos \theta \neq -1$ then they are homotopic 4-forms in $H_+^4(M; \mathbf{R})$, and if Ω_1 is also QK, $p_1(E_0) = p_1(E_1)$.*

In this case a homotopy can be given by $\Omega_t = \cos(t)\Omega_0 + \sin(t)\Omega_2$, where $\Omega_2 = \frac{\Omega_1 - \cos \theta \Omega_0}{\sin \theta}$ is a closed 4-form orthogonal to Ω_0 , and $\Omega_{t=\theta} = \Omega_1$. If Ω_1 is AQK then so it is Ω_t .

Let (J_1, J_2, J_3) be an hyper-Kähler (HK) structure on $\mathbf{R}^8$. This is an o.n. system (of norm $\sqrt{8}$) for the Hilbert-Schmidt inner product. For each $x = (a, b, c) \in \mathbf{R}^3$ let $J_x = aJ_1 + bJ_2 + cJ_3$. The elements with $x \in S^2$ are also called the HK structure of $\mathbf{R}^8$ determined by J_1, J_2 . Let $\mathcal{HK}$ be the set of such HK structures (for a given orientation of $\mathbf{R}^8$). Given an oriented 4-dimensional subspace P of $\mathbf{R}^8$ we take $J'_x, J''_x, x \in S^2$, the HK structures on P and on $P^\perp$ respectively, defined by the g -orthogonal positive complex structures. Then $J_x(X) = J'_x(X^\top) + J''_x(X^\perp)$, where $X^\top$ and $X^\perp$ are respectively the orthogonal projections of X onto P and $P^\perp$, defines a HK structure on $\mathbf{R}^8$. Note that J_x and its fundamental 4-form Ω do not depend on the d.o.n. basis we choose for P and for $P^\perp$. Reciprocally, given (J_1, J_2, J_3) and $X, Y \in \mathbf{R}^8$ unit vectors, with $H_X \perp H_Y$ then $P = H_X = \text{span}\{X, J_1X, J_2X, J_3X\}$, $P^\perp = \text{span}\{Y, J_1Y, J_2Y, J_3Y\} \in \text{Gr}(4, 8)$ and the previous construction w.r.t $P, P^\perp$ recovers $(J_x)_{x \in S^2}$. Let

$$\mathcal{T} = \left\{ \Omega \in \wedge^4(\mathbf{R}^8) : \Omega = \frac{1}{6}(\omega_1 \wedge \omega_1 + \omega_2 \wedge \omega_2 + \omega_3 \wedge \omega_3) : (J_1, J_2, J_3) \in \mathcal{HK} \right\} \subset \wedge_+^4(\mathbf{R}^8)$$

where $\omega_i(u, v) = g(J_i u, v)$. If M is an oriented 8-dimensional manifold then we define the corresponding bundles over M , $\mathcal{HK}(M)$ and $\mathcal{T}(M)$. Now we look for obstructions for two fundamental 4-forms Ω_0 and Ω_1 on M to be homotopic in $\mathcal{T}(M)$. Let us assume that on a neighbourhood U of a point of M , a unit vector field X is defined, and $E_0 = \text{span}\{J_1, J_2, J_3\}$ and $E_1 = \text{span}\{J'_1, J'_2, J'_3\}$ with $J_1 = J'_1$. On what follows we argue as in [4]. At each point, the

subspace $H'_X = \text{span}\{X, J'_1 X, J'_2 X, J'_3 X\}$ is a J_1 -complex subspace of real dimension 4. Since M has dimension 8 there exist a unit vector $Y' \in H'^{\perp}_X \cap H^{\perp}_X$. One can see from (11.2) below that $d = \dim(H_X \cap H'_X) = \dim(H^{\perp}_X \cap H'^{\perp}_X)$. This dimension d is either 2 or 4 (the spaces are J_1 -complex). Thus, on U , if $\text{rank } E_0 \cap E_1$ is constantly equal to one, then d is constantly equal to 2, and we may locally smoothly choose Y' . Set $x = (1, 0, 0)$. Since $TM = H_X \oplus H_{Y'}$, orthogonal sum, there exist unique smooth maps $y, y', u, u' : U \rightarrow \mathbf{R}^3$ such that

$$\begin{aligned} B &= \{X, J_x X, J_y X + J_{y'} Y', J_x(J_y X + J_{y'} Y')\} \\ B^{\perp} &= \{Y', J_x Y', J_u Y' + J_{u'} X, J_x(J_u Y' + J_{u'} X)\}, \end{aligned} \quad (11.2)$$

define a d.o.n. basis of H'_X and $H'_{Y'}$ respectively, where $\|y\|^2 + \|y'\|^2 = \|u\|^2 + \|u'\|^2 = 1$, $g(x, y) = g(x, u) = 0$, $g(x, y') = g(x, u') = 0$, $g(u, y') + g(y, u') = 0$. Away from Q_0 -totally-complex points of H'_X , that is $y(p) \neq 0$, and since Q_0 -quaternionic (and anti- Q_0 -quaternionic) points do not exist as well, i.e. $y'(p) \neq 0$, we have

$$J_y X + J_{y'} Y' = J_{\frac{y}{\|y\|}} (\|y\| X + \|y'\| Y) \quad \text{where } Y = -J_{\frac{y}{\|y\|}} J_{\frac{y'}{\|y'\|}} Y' \in H^{\perp}_X. \quad (11.3)$$

Thus, there exist $x_1 = x, x_2 = \frac{y}{\|y\|}, x_3 = x_1 x_2$ a d.o.n. basis of $\mathbf{R}^3$, and ζ s.t.

$$\begin{aligned} B &= \{X, J_{x_1} X, J_{x_2}(cX + sY), J_{x_3}(cX + sY)\} \\ B^{\perp} &= \{Y, J_{x_1} Y, J_{x_2}(cY - sX), J_{x_3}(cY - sX)\}, \end{aligned} \quad (11.4)$$

where $c = \cos \zeta > 0$, $s = \sin \zeta > 0$, define a d.o.n. basis of H'_X and $H'_{Y'}$ respectively. Furthermore $Y \in H^{\perp}_X \cap H'^{\perp}_X$ as well. Note that if we make $y \rightarrow 0$, we are approaching a Q_0 -totally-complex point, we have at $y(p) = 0$, $J_y X + J_{y'} Y' = J_{y'} Y'$, but we may not smoothly or continuously extended x_2 and Y as $x_2 = y'$ and $Y = Y'$, or smoothly extend c to 0 and s to 1 at those points. This ambiguity will imply an ambiguity on the choice of $e_3, \dots, e_8$ bellow, and we need to use (11.4) to define the homotopy in $\mathcal{T}(M)$. A similar problem occurs at Q_0 -quaternionic points of H'_X , where $H_X \cap H'_X$ becomes 4-dimensional. A smooth choice of Y' might be impossible. Set $e_1 = X$, $e_2 = J_{x_1} X$, $e_3 = J_{x_2} X$, $e_4 = J_{x_3} X$, $e_5 = Y$, $e_6 = J_{x_1} Y$, $e_7 = J_{x_2} Y$, $e_8 = J_{x_3} Y$. Then $\omega_1 = "J_{x_1}" = e^{12} + e^{34} + e^{56} + e^{78}$, $\omega_2 = "J_{x_2}" = e^{13} - e^{24} + e^{57} - e^{68}$, $\omega_3 = "J_{x_3}" = e^{14} + e^{23} + e^{58} + e^{67}$. We define $\omega_4 = "J_4" = e^{17} - e^{28} + e^{35} - e^{46}$, $\omega_5 = "J_5" = e^{18} + e^{27} + e^{36} + e^{45}$. Then we have $\omega'_1 = \omega_1$, $\omega'_2 = c\omega_2 + s\omega_4$, $\omega'_3 = c\omega_3 + s\omega_5$, and so locally,

$$E_1 = (J_{x_1}, cJ_{x_2} + sJ_4, cJ_{x_3} + sJ_5) \quad (11.5)$$

$$\Omega_1 = c^2 \Omega_0 + \frac{1}{3} s c (\omega_2 \wedge \omega_4 + \omega_3 \wedge \omega_5) + \frac{s^2}{6} (\omega_1 \wedge \omega_1 + \omega_4 \wedge \omega_4 + \omega_5 \wedge \omega_5) \quad (11.6)$$

$$\tilde{H} = \frac{s^2}{6} ((\omega_1 \wedge \omega_1 + \omega_4 \wedge \omega_4 + \omega_5 \wedge \omega_5) - \frac{7}{15} (\omega_1 \wedge \omega_1 + \omega_2 \wedge \omega_2 + \omega_3 \wedge \omega_3)) + \frac{2}{3} s c (\omega_2 \wedge \omega_4). \quad (11.7)$$

We observe that $J_{x_1}, J_{x_2}, J_{x_3}, J_4, J_5$ is an o.n. system in $\wedge^2 TM$, and J_{x_1} anti-commutes with $J_{x_2}, J_{x_3}, J_4, J_5$, and $J_{x_3} = J_{x_1} J_{x_2}$, $J_5 = J_{x_1} J_4$, J_{x_2} anti-commutes with J_4, J_{x_3} anti-commutes with J_5 . This explains that (11.5) defines an HK structure, and it is elementary to verify it spans E_1 . We also have $\omega_2 \wedge \omega_4 = \omega_3 \wedge \omega_5$ and $\cos \theta = \frac{1}{15}(7 + 8c^2) \in [\frac{7}{15}, 1]$ is positive, (and c can be continuously extended to Q_0 -quaternionic points and to Q_0 -totally-complex points). Thus $\cos \theta \neq -1$ everywhere. So we may conclude:

Proposition 11.3 *If Q_0 and Q_1 are two almost quaternionic Hermitian structures on a 8-manifold M with $\text{rank}(E_0 \cap E_1) \geq 1$, then Ω_1 and Ω_0 are homotopic in $\Lambda_+^4(TM)$. If $E_0 \cap E_1$ has constant rank one and $\cos \theta > \frac{7}{15}$, then Ω_0 and Ω_1 are locally homotopic in $\mathcal{T}(M)$.*

Proof. Locally we may take for each $t \in [0, 1]$ B_t and $B_t^\perp$ as in (11.4) replacing c and s by $c(t) = \cos(t\zeta)$, $s(t) = \sin(t\zeta)$. Then $\Omega_t = \frac{1}{6}(\omega_1(t) \wedge \omega_1(t) + \omega_2(t) \wedge \omega_2(t) + \omega_3(t) \wedge \omega_3(t)) \in \mathcal{T}(M)$ where $J_1(t) = J_{x_2}$, $J_2(t) = c(t)J_{x_2} + s(t)J_4$, $J_3(t) = J_1(t)J_2(t)$. QED

Corollary 11.1 *If Q_0 and Q_1 are both QK, then one of the three cases holds: (a) $E_0 = E_1$, (b) $E_0 \cap E_1$ has rank one and both Q_0, Q_1 are locally HK structures, and M is Kähler. Furthermore $p_1(E_0) = p_1(E_1)$. (c) $E_0 \cap E_1 = \{0\}$.*

Proof. $E_0 \cap E_1$ is a parallel subbundle of $\Lambda^2(TM)$, and so it has constant rank. If it has rank ≥ 2 then it must have rank 3, and so $E_0 = E_1$. So let us assume it has rank 1. This means that both E_0 and E_1 have a common compatible parallel complex structure. This implies M is Kähler and E_0 and E_1 are locally HK (see [3]). QED

Remark 7. If we want to find a homotopy by closed 4-forms and Q_0 or Q_1 are not AQK we have to compute $p_1(E_0)$ and $p_1(E_1)$ directly from local frames (J_1, J_2, J_3) and (J'_1, J'_2, J'_3) . Assume we are in the simplest case that Q_0 is HK, $E_0 \cap E_1$ has rank ≥ 1 and Λ_+^4 is flat. We may compare $p_1(E_1)$ with $p_1(E_0) = 0$ using a homotopy $p_1(E_t)$, locally defined by $B_t, B_t^\perp$ as in the previous proofs, and using the bundle $\tilde{E}$ over $M \times [0, 1]$, by computing $p_1(\tilde{E})(X, \frac{d}{dt})$ as we did in section 9, using the remark 4 of section 7. This would lead to a transgression form with singularities at Q_0 -totally complex points and at Q_0 -quaternionic points, and so to a residue formula. This is difficult, because B_t are only locally defined, and it requires more investigation on Λ_+^4 and $\mathcal{T}$, namely, under a musical isomorphism, by considering the elements of $\mathcal{T}$ as isomorphic and symmetric operators of $\Lambda^2 TM$, whose eigenvectors are almost complex structures. So we will not discuss this problem in this paper and leave it for later.

References

- [1] D.V. Alekseevsky, *Quaternion Riemannian spaces with transitive reductive or solvable group of motions*, Funktional Anal. i Prilozhen. **4** (1970) 68-69.
- [2] D.V. Alekseevsky & S. Marchiafava, *Almost quaternionic Hermitian and Quasi-Kähler manifolds*. Proc. of the international workshop on almost complex structures, K. Sekigawa et al. (ed.), August 20-25, 1992. Singapore: World Scientific. 150-175 (1994)
- [3] D.V. Alekseevsky & S. Marchiafava & M. Pontecorvo, *Some geometric aspects of compatible complex structures on a quaternionic Kähler manifold*. Differential Geometry and Applications, Satellite Conf. of ICM in Berlin, Aug.10-14, 1998, Brno Masaryk Univ. in Brno (Czech Republic) 1999, 1-11.
- [4] D.V. Alekseevsky & A. Pereira do Vale & I. Salavessa, *Minimal submanifolds of quaternionic Kähler manifolds*, in preparation.
- [5] V. Apostolov & T. Draghici, *The curvature and the integrability of almost-Kähler manifolds: a survey*. Symplectic and contact topology: interactions and perspectives (Toronto/ ON/Montreal, QC, 2001), 25-53, Fids Inst. Commun., 35, Amer. Math. Soc., Providence, RI, 2003.

- [6] M. Atiyah & R. Bott & V.K. Patodi, *On the heat equation and the index theorem*, Invent. Math. **19** (1973), 279-330.
- [7] A.L. Besse, *Géométrie riemannienne en dimension 4*, Cedric-Fernand Nathan, Paris (1981).
- [8] A.L. Besse, *Einstein manifolds*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3). Springer-Verlag, Berlin, 1987.
- [9] D.E. Blair, *The "total scalar curvature" as a symplectic invariant and related results*, Proceedings of the 3rd Congress of Geometry, Thessaloniki/Greece 1991, (1992), 79-83.
- [10] F. Costantino, *Branched shadows on complex structures on 4-manifolds*, math.GT/0502292.
- [11] J-H. Eschenburg & I.V. Guadalupe & R.A. Tribuzy, *The Fundamental Equations of Minimal Surfaces in CP^2* , Math. Ann. **270** (1985) 571-598.
- [12] S. Gallot & D. Hulin & J. Lafontaine, *Riemannian Geometry*, Springer-Verlag, Universitext 1990.
- [13] P. Gauduchon, *Hermitian connections and Dirac operators*, Boll. U.M.I.(7) 11-B (1997), suppl. fasc. 2, 257-288
- [14] A. Gray, *Tubes*, 2nd edition, Progress in Mathematics **221**, Birkhäuser Verlag.
- [15] R. Harvey & H.B. Lawson, Jr. *Geometric Residue Theorems*, Amer. J.Math. **117** (1995) (4), 829-873.
- [16] R. Harvey & H. B. Lawson, Jr. *A theory of Characteristic currents associated with singular connection*, Bull. Amer. Math. Soc. (NS) **31** (1994) 54-63.
- [17] R. Harvey & S.W. Semmes, *Zero divisors of atomic functions*, Ann. of Math. (2) **135**(3) (1992), 567-600.
- [18] N.J. Hitchin, *The self-duality equations on a Riemann surface*, Proc. Lond. Math. Soc. (3) **55**(1987), 59-126.
- [19] D. Joyce, *Compact manifolds with special holonomy*, Oxford Mathematical Monographs series, 2000.
- [20] V.Y. Kraines, *Topology of quaternionic manifolds*, Trans. AMS **122**, 357-367 (1966).
- [21] M.J. Micallef & J.D. Moore *Minimal two-spheres and the topology of manifolds with positive curvature on totally isotropic two-planes*, Annals of Math. **127** (1988), 199-227.
- [22] S.M. Salamon, *Almost parallel structures*, in *Global differential geometry: the mathematical legacy of Alfred Gray*, (Bilbao 2000), Contemp. Math. **288**. Amer. Math. Soc., Providence R.I. 2001, 162-181.
- [23] I. Salavessa & A. Pereira do Vale, *Cayley submanifolds of Calabi-Yau 4-folds*, math.DG/0408206
- [24] I. Salavessa & G. Valli, *Minimal submanifolds of Kähler-Einstein manifolds with equal Kähler angles*, Pacific J. Math. **205**(1)(2002), 197-235.
- [25] I.M. Singer & J.A. Thorpe, *The curvature of 4-dimensional Einstein spaces*, Global Analysis (Papers in Honor of K. Kodaira) 355-365, Univ. Tokyo Press, Tokyo.
- [26] S.M. Webster, *Minimal surfaces in a Kähler surface*, J. Diff. Geo. **20** (1984), 463-470.
- [27] J.G. Wolfson, *Minimal Surfaces in Kähler Surfaces and Ricci Curvature*, J. Diff. Geo. **29** (1989), 281-294.